

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
12 December 2002 (12.12.2002)

PCT

(10) International Publication Number
WO 02/099079 A2

(51) International Patent Classification⁷: C12N

(74) Agent: DECAMP, James, D., Clark & Elbing LLP, 101
Federal Street, Boston, MA 02110 (US).

(21) International Application Number: PCT/US02/18066

(22) International Filing Date: 6 June 2002 (06.06.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/296,554 6 June 2001 (06.06.2001) US

(63) Related by continuation (CON) or continuation-in-part
(CIP) to earlier application:
US 60/296,554 (CIP)
Filed on 6 June 2001 (06.06.2001)

(71) Applicant (for all designated States except US): **THE
GENERAL HOSPITAL CORPORATION** [US/US]; 55
Fruit Street, Boston, MA 02114 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **SHEEN, Jen**
[US/US]; 9 Hawthorne Place, Apartment 5J, Boston, MA
02114 (US). **HWANG, Ildoo** [KR/US]; 29J Montrose
Manor Court, Baltimore, MD 21228 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

Published:

*without international search report and to be republished
upon receipt of that report*

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: CYTOKININ RESPONSE REGULATORS AND USES THEREOF

(57) Abstract: The invention generally features methods for increasing yield, shoot formation, and delaying senescence in plants with the use of transgenes that regulate the cytokinin response. The invention also features plants and plant components that harbor the transgene(s).

WO 02/099079 A2

CYTOKININ RESPONSE REGULATORS AND USES THEREOF

5

Background of the Invention

This invention relates to genetically-engineered plants having increased yield and productivity, shoot, leaf and meristem formation, enhanced photosynthesis, and delayed senescence.

10 Despite long recognition of cytokinins as essential plant hormones involved in diverse processes of plant growth and development, including cell division, shoot initiation, leaf and root differentiation, chloroplast biogenesis, apical dominance, and senescence, the molecular and biochemical mechanisms underlying cytokinin actions have not been elucidated (Davies, *Plant Hormones: Physiology, Biochemistry and Molecular Biology*, 1995, Kluwer Academic Publishers, Dordrecht; Meijer and Murray, *Curr. Opin. Plant Biol.* 4: 44-9, 2001; Mok and Mok, *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 52: 89-118, 2001; Quirino et al., *Trends Plant Sci.* 5: 278-82, 2000). Recent genetic identification of *Arabidopsis* hybrid histidine protein kinases (AHKs), CKI1 and CRE1, in
15 cytokinin signalling (Kakimoto, *Science* 274: 982-5, 1996; Suzuki et al., *Plant Cell. Physiol.* 42: 107-13, 2001; Inoue et al., *Nature* 409: 1060-3, 2001), the characterization of *Arabidopsis* response regulators (ARRs) as cytokinin primary response genes (D'Agostino et al., *Plant Physiol.* 124: 1706-17, 2000; Kiba et al., *Plant Cell Physiol.* 40: 767-71, 1999), and the existence of *Arabidopsis* histidine-
20 containing phosphotransmitters (AHPs; Suzuki, Imamura, et al., *Plant Cell Physiol.* 39: 1258-68, 1998), have implicated the involvement of a two-component phosphorelay mechanism in cytokinin signal transduction. However, cellular and molecular evidence is lacking in supporting a complete cytokinin signalling circuit in plant cells.

30 Two-component circuitry, consisting of a histidine kinase (HK) sensor and a response regulator (RR) output, are responsible for signal transduction in most prokaryotic and some eukaryotic systems. The signalling pathway is initiated by

a HK sensor and mediated by phosphotransfer between a conserved histidine (His) residue in HKs or histidine-containing phosphotransmitters (HPs) and a conserved aspartate (Asp) residue in RRs (Wurgler-Murphy and Saito, *Trends Biochem. Sci.* 22: 172-6, 1997; Stock et al., *Annu. Rev. Biochem.* 69: 183-215, 5 2000). Since there are only one HK, one HP, and two RRs in *S. cerevisiae*, it was previously believed that two-component signal transduction has limited function in eukaryotes (Stock et al., *supra*). However, the completion of the *Arabidopsis* genome has revealed over 40 genes encoding putative two-component signal transducers, AHKs, AHPs and ARRs, suggesting a significant involvement of the 10 ancient and conserved signalling mechanism in many facets of plant cell regulation (Urao, et al., *Trends Plant Sci.* 5: 67-74, 2000). The identification of conserved HK signature motifs and/or activity in the photoreceptor phytochrome, putative osmosensor, and the ethylene and cytokinin receptors in *Arabidopsis* further supports this view (Inoue et al., *supra*; Urao, et al., *supra*; Yeh and 15 Lagarias, *Proc. Natl. Acad. Sci. U SA* 95: 13976-81, 1998; Bleecker and Kende, *Annu. Rev. Cell Dev. Biol.* 16: 1-18, 2000).

In eukaryotic signal transduction, the two-component circuit often provides a link between the HK sensor to a MAP kinase (MAPK) signaling cascade. For example, the osmosensing signal transduction pathway in yeast is 20 mediated by the SLN1/YPD1/SSK1 phosphorelay. The RR SSK1 then activates the HOG1 MAPK cascade in the cytosol to control gene expression (Maeda et al., *Nature* 369: 242-5, 1994; Posas et al., *Cell* 86: 865-75, 1996; Posas and Saito, *EMBO J.* 17: 1385-94, 1998). It has been speculated that plant HKs such as the ethylene receptor (ETR1) and a putative osmosensor (AHK1) can initiate a 25 phosphorelay and transmit the signal through a MAPK cascade (Bleecker and Kende, *supra*; Gamble et al., *Proc. Natl. Acad. Sci. U SA* 95: 7825-9, 1998; Clark et al., *Proc. Natl. Acad. Sci. U SA* 95: 5401-6, 1998; Urao et al., *Plant Cell* 11: 1743-54, 1999). Due to the lack of physiological plant cell assays, the mechanisms of HK action and the signal transduction pathways of any HK- 30 mediated plant responses, including cytokinin signalling, remain obscure.

Summary of the Invention

The present invention capitalizes on the discovery that manipulation of the expression of cytokinin response regulators increases plant yield growth, and
5 productivity.

In one aspect, the invention features a method for increasing yield in a plant, the method including the steps of: (a) introducing into plant cells a transgene including DNA encoding a B-type response regulator operably linked to a promoter functional in plant cells to yield transformed plant cells; and (b)
10 regenerating a plant from the transformed cells, wherein the B-type response regulator is expressed in the cells of the transgenic plant, thereby increasing yield in the plant. In preferred embodiments, the B-type response regulator is a crucifer B-type response regulator (for example, ARR1 (SEQ ID NO.: 2); ARR2 (SEQ ID NO.: 3); and ARR10 (SEQ ID NO.: 9)).

15 In a second aspect, the invention features a method for increasing yield in a plant, the method including the steps of: (a) introducing into plant cells a transgene operably linked to a promoter functional in plant cells to yield transformed plant cells; and (b) regenerating a plant from the transformed cells, wherein expression of the transgene reduces expression of an A-type response
20 regulator in the cells of the plant, thereby increasing yield in the plant. In preferred embodiments, the A-type response regulator is a crucifer A-type response regulator (for example, ARR 4 (SEQ ID NO.: 5); ARR 5 (SEQ ID NO.: 6); ARR 6 (SEQ ID NO.: 7); and ARR 7 (SEQ ID NO.: 8)). In other preferred embodiments, the transgene expresses an antisense molecule of A-type response
25 regulator; a dominant negative gene product of A-type response regulator; or expression of the transgene results in the co-suppression of the A-type response regulator.

In a third aspect, the invention features a method for increasing shoot formation in a plant, the method including the steps of: (a) introducing into plant
30 cells a transgene including DNA encoding a B-type response regulator operably

linked to a promoter functional in plant cells to yield transformed plant cells; and
(b) regenerating a plant from the transformed cells, wherein the B-type response
regulator is expressed in the cells of the plant, thereby increasing shoot formation
in the plant. In preferred embodiments, the method includes the use of a B-type
5 response regulator that is a crucifer B-type response regulator.

In a fourth aspect, the invention features a method for increasing shoot
formation in a plant, the method including the steps of: (a) introducing into plant
cells a transgene operably linked to a promoter functional in plant cells to yield
transformed plant cells; and (b) regenerating a plant from the transformed cells,
10 wherein expression of the transgene reduces expression of an A-type response
regulator in the cells of the plant, thereby increasing shoot formation in the plant.

In a fifth aspect, the invention features a method for delaying senescence
in a plant, the method including the steps of: (a) introducing into plant cells a
transgene including DNA encoding a B-type response regulator operably linked
15 to a promoter functional in plant cells to yield transformed plant cells; and (b)
regenerating a plant from the transformed cells, wherein the B-type response
regulator is expressed in the cells of the plant, thereby delaying senescence in the
plant.

In a sixth aspect, a method for delaying senescence in a plant, the method
20 including the steps of: (a) introducing into plant cells a transgene operably linked
to a promoter functional in plant cells to yield transformed plant cells; and (b)
regenerating a plant from the transformed cells, wherein expression of the
transgene reduces expression of an A-type response regulator in the cells of the
plant, thereby delaying senescence in the plant.

25 In a seventh aspect, the invention features a method for increasing yield in
a plant, the method including the steps of: (a) introducing into plant cells a
transgene including DNA encoding a histidine kinase operably linked to a
promoter functional in plant cells to yield transformed plant cells; and (b)
regenerating a plant from the transformed cells, wherein histidine kinase is

30

expressed in the cells of the plant, thereby increasing yield in the plant. In preferred embodiments, the histidine kinase is a crucifer histidine kinase (for example, CKI1 (SEQ ID NO.: 13) and CRE1 (SEQ ID NO.: 18)).

In an eighth aspect, the invention features a method for increasing shoot formation in a plant, the method including the steps of: (a) introducing into plant cells a transgene including DNA encoding a histidine kinase operably linked to a promoter functional in plant cells to yield transformed plant cells; and (b) regenerating a plant from the transformed cells, wherein histidine kinase is expressed in the cells of the plant, thereby increasing shoot formation in the plant.

In a ninth aspect, a method for delaying senescence in a plant, the method including the steps of: (a) introducing into plant cells a transgene including DNA encoding a histidine kinase operably linked to a promoter functional in plant cells to yield transformed plant cells; and (b) regenerating a plant from the transformed cells, wherein histidine kinase is expressed in the cells of the plant, thereby delaying senescence in the plant.

The invention further includes plants and plant components expressing one, or a combination of two or more, of the aforementioned transgenes. The invention also encompasses methods of generating these plants or plant components, which may involve introducing the transgenes individually to separate plants or plant components, and then crossing the appropriate genotypes to get the desired transgene combination. Alternatively, one or more constructs expressing a desired combination of transgenes may be generated and then introduced into the plant or plant component.

For example, the invention also relates to a plant or plant component comprising at least one transgene encoding (i) an A-type response regulator polypeptide, (ii) an antisense A-type response regulator RNA, (iii) a dominant negative A-type response regulator polypeptide, (iv) a B-type response regulator polypeptide, (v) an antisense B-type response regulator RNA, (vi) a dominant negative B-type response regulator, (vii) an antisense HK RNA, (viii) a dominant

negative HK polypeptide, or any combination of (i) –(viii). In addition, the invention includes any of the aforementioned plants further including a transgene encoding a histidine kinase polypeptide.

The invention further includes a plant or plant component that includes:

- 5 (a) a first transgene encoding (i) an A-type response regulator polypeptide, (ii) an antisense A-type response regulator RNA, or (iii) a dominant negative A-type response regulator polypeptide or combination thereof; (b) a second transgene encoding (iv) a B-type response regulator polypeptide, (v) an antisense B-type response regulator RNA, (vi) a dominant negative B-type response regulator or
10 combination thereof, and (c) a third transgene encoding (vii) a HK polypeptide; (viii) an antisense HK RNA, (ix) a dominant negative HK polypeptide or combination thereof.

Exemplary plants which are useful in the methods of the invention, as well as for generating the plants (or plant cells, plant components, plant tissues, or
15 plant organs) of the invention, include dicots and monocots, such as sugar cane, wheat, rice, maize, sugar beet, barley, manioc, crucifer, mustard, potato, soybean, sorghum, cassava, banana, grape, oats, tomato, millet, coconut, orange, rye, cabbage, apple, eggplant, watermelon, canola, cotton, carrot, garlic, onion, pepper, strawberry, yam, papaya, peanut, onion, legume, bean, pea, mango, and
20 sunflower.

By “operably linked” is meant that a gene and a regulatory sequence(s) are connected in such a way as to permit gene expression when the appropriate molecules (for example, transcriptional activator proteins) are bound to the regulatory sequence(s).

25 By “plant cell” is meant any self-propagating cell bounded by a semi-plant component expression control permeable membrane and containing a plastid. Such a cell also requires a cell wall if further propagation is desired. Plant cell, as used herein, includes, without limitation, algae, cyanobacteria, seeds, suspension cultures, embryos, meristematic regions, callus tissue, leaves, roots, shoots,
30 gametophytes, sporophytes, pollen, and microspores.

By "plant component" is meant a part, segment, or organ obtained from an intact plant or plant cell. Exemplary plant components include, without limitation, somatic embryos, leaves, stems, roots, flowers, tendrils, fruits, scions, and rootstocks.

5 By "transgene" is meant any piece of DNA which is inserted by artifice into a cell, and becomes part of the genome of the organism which develops from that cell. Such a transgene may include a gene which is partly or entirely heterologous (i.e., foreign) to the transgenic organism, or may represent a gene homologous to an endogenous gene of the organism.

10 By "transgenic" is meant any cell which includes a nucleic acid sequence (e.g., a recombinant DNA sequence) which is inserted by artifice into a cell and becomes part of the genome of the organism which develops from that cell. As used herein, the transgenic organisms are generally transgenic plants and the DNA (transgene) is inserted by artifice into the nuclear or plastidic genome.

15 By "yield" or "plant yield" is meant increased growth (e.g., crop growth) or increased biomass. For example, increased yield results from increased shoot growth or meristem formation. Plants expressing the genes disclosed herein exhibiting increased yield can be selected by visual observation, for example by comparison with a wild-type plant.

20 By "reducing expression" or "reduces expression" is meant a decrease in the level of gene expression (for example, expression of a gene encoding a A-type response regulator) by at least 30-50%, preferably by at least 50-80%, and more preferably by at least 80-95% or greater relative to the level in a control plant (for example, a wild-type plant). Reduction of such expression levels may be
25 accomplished by employing standard methods which are known in the art including, without limitation, antisense and co-suppression technologies, expression of a dominant negative gene product, or through the generation of mutated genes using standard mutagenesis techniques. Levels of negative

regulator polypeptide or transcript are monitored according to any standard technique including, but not limited to, northern blotting, RNase protection, or immunoblotting.

By "crucifer" is meant any plant that is classified within the Cruciferae family. The Cruciferae include many agricultural crops, including, without
5 limitation, rape (for example, *Brassica campestris* and *Brassica napus*), broccoli, cabbage, brussel sprouts, radish, kale, Chinese kale, kohlrabi, cauliflower, turnip, rutabaga, mustard, horseradish, and *Arabidopsis*.

By "a promoter functional in a plant cell" is meant any minimal sequence
10 sufficient to direct transcription in a plant cell. Included in the invention are promoter elements that are sufficient to render promoter-dependent gene expression controllable for cell-, tissue-, or organ-specific gene expression, or elements that are inducible by external or internal agents (for example, cytokinins), or elements that are capable of cycling gene transcription; such
15 elements may be located in the 5' or 3' regions of the native gene or engineered into a transgene construct.

The invention provides a number of important advances and advantages for improving and enhancing agronomically important traits such as photosynthesis, productivity, yield, leaf, shoot, and meristem formation, and
20 delaying senescence. In particular, the invention provides for increased production efficiency, as well as for improvements in quality and yield of crop plants and ornamentals. Thus, the invention contributes to the production of high quality and high yield agricultural products; for example, fruits, ornamentals, vegetables, cereals, and field crops. Genetically-improved seeds and other plant
25 products that are produced using plants expressing the genes and methods described herein also render farming possible in areas previously unsuitable for agricultural production. The mechanisms disclosed herein for increasing plant yield and productivity is expected to be ubiquitous throughout the plant kingdom.

Other features and advantages of the invention will be apparent from the
30 following description of the preferred embodiments thereof, and from the claims.

Brief Description of the Drawings

Figure 1A is a bar graph showing the specificity of plant hormone responses in the *Arabidopsis* mesophyll protoplast transient expression system.

5 **Figure 1B** is a graph depicting a cytokinin dose response for the *ARR6* promoter induction.

Figure 1C is a bar graph showing the induction of the *ARR6* promoter by different cytokinins.

Figure 1D shows that CKI1 activation of *ARR6-LUC* requires HK activity and phosphorelay. The photograph (top) shows an autoradiogram depicting the expression of CKI1 proteins. The bar graph (bottom) shows the results of transfected protoplasts that were incubated for 3 hours to allow effector gene expression before cytokinin treatment for another 3 hours.

15 **Figure 1E** shows that CRE1 confers cytokinin hypersensitivity on the activation of the *ARR6-LUC* reporter. The photograph (top) shows an autoradiogram depicting the expression of *Arabidopsis* HK proteins. The results of this incubation with cytokinin are shown in the bar graph (bottom).

Figure 1F shows a series of photomicrographs showing that CKI1 is localized at the plasma membrane.

20 **Figure 2A** demonstrates that AHP acts as a shuttle between the cytosol and nucleus in cytokinin signalling. The photograph (top) depicts an autoradiogram demonstrating expression of AHP proteins. The bar graph (bottom) shows the effect of cytokinin treatment on plasmid expression.

Figure 2B shows a series of photomicrographs demonstrating that
25 cytokinin induces transient AHP translocation.

Figure 3A shows negative and positive regulation by ARRs. The photograph depicts an autoradiogram (top) showing the expression of *Arabidopsis* ARR proteins. The bar graph (bottom) shows the activity of *ARR6* promoter activity when cells were then incubated with or without 100 nM *t*-zeatin
30 for another 3 hours before the activity was measured.

Figure 3B shows that ARR2s act downstream of CKI1 in cytokinin signalling.

Figure 3C shows a series of photomicrographs demonstrating that *Arabidopsis* ARR2s are localized in the nucleus.

5 **Figure 4A** is a photograph showing that ectopic expression of ARR2 is sufficient to promote cytokinin responses in transgenic tissues and plants when exposed to auxin.

10 **Figure 4B** is a photograph showing that ectopic expression of ARR2 is sufficient to promote cytokinin responses in transgenic tissues and plants when exposed to both auxin and cytokinin.

Figure 4C is a photograph demonstrating that cytokinin signaling delays dark-induced senescence in transgenic *Arabidopsis* plants.

Figure 5 is a schematic of a model of the cytokinin signal transduction pathway in *Arabidopsis*.

15

Detailed Description

Overview

20 Cytokinins are essential plant hormones involved in shoot meristem and leaf formation, cell division, chloroplast biogenesis, and senescence. Although hybrid histidine protein kinases (HKs) have been implicated in cytokinin perception in *Arabidopsis*, the action of HK receptors and downstream signalling pathways remain elusive in plant cells. To study the action of HK receptors and downstream signaling pathways, a mesophyll protoplast transient assay based on the transcription activation of a cytokinin primary response gene *ARR6* in
25 *Arabidopsis* was developed. The use of this high-throughput cell assay for functional genomic analysis of the two-component regulators has provided a powerful tool to decipher the first complete cytokinin signalling pathway. Rather than following the established eukaryotic HK and MAPK cascade paradigm,

plant two-component signalling can integrate multiple HK activities to common AHPs as cytoplasm and nuclear shuttles, and distinct nuclear ARR_s as master regulatory outputs.

As is described in more detail below, a new eukaryotic two-component signalling circuit that initiates cytokinin signalling by employing distinct hybrid HK activities at the plasma membrane has been identified. In particular, AHPs are shown to act as novel eukaryotic signalling shuttles between the cytoplasm and nucleus in a transient and cytokinin-dependent manner. The short signalling circuit reaches the nuclear target genes by de-repressing nuclear response regulators ARR₁, ARR₂, and ARR₁₀ as transcription activators. The cytokinin-inducible ARR₄, ARR₅, ARR₆, and ARR₇ act as transcription repressors, suggesting a negative feedback loop. Thus, the *Arabidopsis* cytokinin signal transduction pathway consists of four major steps: HK sensing and signalling, AHP translocation, ARR de-repression, and a negative feedback loop by cytokinin-inducible ARR_s. Transgenic tissue and plant analyses support the importance of this core signalling pathway in diverse cytokinin responses. Furthermore, ectopic expression of the master regulator ARR₂ as the rate-limiting two-component output in transgenic *Arabidopsis* was sufficient to mimic cytokinin in promoting shoot meristem proliferation, leaf differentiation, expansion, and longevity.

The examples provided below are for the purposes of illustrating the invention, and should not be construed as limiting.

Cytokinin Signal is Perceived and Initiated by the Multiple HK receptors

To elucidate the regulatory circuitry in cytokinin signal transduction, a leaf cell assay based on cytokinin inducible transcription in *Arabidopsis* mesophyll protoplasts was developed. The 2.4 kb promoter of an *Arabidopsis* cytokinin primary response gene (SEQ ID NO.: 1), encoding the response regulator 6 (ARR₆), was fused to the firefly luciferase (LUC) reporter. In Fig. 1A, transfected protoplasts were incubated with or without 100 nM *t*-zeatin, 1 μM

indole-3-acetic acid (IAA), or 100 μ M ABA. After 3 hours of incubation, LUC and GUS activities were determined. In transfected protoplasts, the activity of *ARR6-LUC* was specifically induced by cytokinin but not by other plant hormones such as abscisic acid (ABA) or auxin (Fig. 1A).

5 In the same system, the *GH3* promoter was only activated by auxin, whereas the *RD29A* promoter was solely induced by abscisic acid (ABA), demonstrating the specificity of three plant hormone responses in *Arabidopsis* mesophyll protoplasts (Fig. 1A). The activity of *ARR6-LUC* was induced by physiological concentration of a natural cytokinin *trans*-zeatin (*t*-zeatin) from 1 to
10 100 nM (Fig. 1B).

To further show the specificity of the cytokinin response, various active and inactive cytokinin analogues were examined. Only the active cytokinins, *t*-zeatin, 2-isopentenyladenine (2-IP; 100 nM) and benzyladenine (BA; 100 nM), induced the reporter gene *ARR6-LUC* (Fig. 1C).

15 The response in the protoplast system is similar to the cytokinin activation of various *ARR* genes shown *in planta* (D'Agostino et al., *supra*; Kiba et al., *supra*). Thus, we have established a reliable and physiological system to dissect the regulatory components in the cytokinin signal transduction pathway in *Arabidopsis*.

20

Cytokinin Signalling by Distinct HK Receptors and Phosphorelay

Despite the genetic identification of histidine kinases (HKs) as receptors for the plant hormones ethylene and cytokinin in *Arabidopsis*, whether or where HK activity is important for signal transduction in plant cells remains an
25 unresolved question due to the lack of physiological plant cell assays. CKI1, a hybrid HK with a conserved response regulator (RR) receiver domain has been implicated in cytokinin responses (Kakimoto, *supra*). The enhanced CKI1 expression in *Arabidopsis shooty callus* mutants also supports its role in cytokinin signalling (Frank et al., *Plant Physiol.* 122: 721-9, 2000). However, the
30 biochemical mechanism of CKI1 function and its role in cytokinin signalling are

still unknown. To determine whether CKI1 mediates cytokinin responsive transcription, the full-length *CKI1* gene (SEQ ID NO.: 13) was tagged with double haemagglutinin epitope (DHA) sequence and fused to a constitutive promoter, and co-transfected with *ARR6-LUC* into the *Arabidopsis* protoplasts.

- 5 Specifically, protoplasts were cotransfected with the *ARR6-LUC* reporter and the effector plasmid expressing CKI1 (SEQ ID NO.: 13), CKI1(H405Q) (SEQ ID NO.: 14), CKI1(D1050N) (SEQ ID NO.: 15), CKI1-GFP, or mER7-GFP (control). As shown in Fig. 1D, the wild-type CKI1 activated the *ARR6* promoter without exogenous *t*-zeatin. It is possible that over-expression of CKI1 renders
- 10 protoplasts hypersensitive to endogenous cytokinin and/or exceeds the capacity of negative regulators. Alternatively, CKI1 could encode a constitutively active HK connected to the cytokinin signal transduction pathway. Cytokinin treatment hardly enhanced the reporter gene activity in the presence of CKI1 (Fig. 1D).

- To determine whether the HK activity and phosphorelay are required for
- 15 CKI1 activation of the *ARR6* promoter, the conserved His 405 and Asp 1040 residues in the HK and phosphorylation sites, respectively, were mutated. Specifically, protoplasts were transfected with plasmid DNA expressing CKI1 (SEQ ID NO.: 13), CKI (H405Q) (SEQ ID NO.: 14) and CKI1(D1040N) (SEQ ID NO.: 15), or vector DNA as a control. The [³⁵S]methionine labeled protein
- 20 from transfected protoplasts was immunoprecipitated with anti-HA and proteinA-agarose, and analyzed by SDS-PAGE. Despite comparable expression of the CKI1(H405Q) and CKI1(D1050N) mutants as the wild-type CKI1, neither could activate the expression of *ARR6-LUC* (Fig. 1D). Furthermore, the CKI1(H405Q) mutant served as a strong dominant negative mutant to block the activation of the
- 25 *ARR6* promoter by exogenous *t*-zeatin. The CKI1(D1040N) mutant is less potent. These results indicated that HK activity and phosphorelay from His 405 to D1040 are required for the CKI1 function in activating cytokinin signalling. Moreover, the dominant negative CKI1(H405Q) mutant might interfere with cytokinin perception and/or disturb downstream signalling.

Recently, another *Arabidopsis* hybrid HK, CRE1/AHK4/WOL, has been shown to be a cytokinin receptor. The evidence is based on the inability of the *cre1* mutant to respond to cytokinin in the shoot induction assay and its ability to complement HK mutants of budding and fission yeast and *E. coli* in a cytokinin-dependent manner (Suzuki et al., *supra*; Inoue et al., *supra*; Mahonen et al., *Genes Dev.* 14: 2938-43, 2000). Since CRE1/AHK4/WOL is predominantly expressed in roots (Mahonen et al., *supra*; Ueguchi et al., *Plant Cell Physiol.* 42: 231-5, 2001) and the *cre1* and *wol* recessive mutants lack an obvious leaf phenotype (Inoue et al., *supra*; Mahonen et al., *supra*; Ueguchi et al., *supra*), it is possible that other closely related *Arabidopsis* HK genes, such as AHK2 and AHK3, and CKII provide cytokinin receptor functions in promoting shoot meristem and leaf formation, and delaying leaf senescence (Inoue et al., *supra*; Ueguchi et al., *supra*).

To directly test the function of these HKs in cytokinin signalling, constructs expressing AHK2 (SEQ ID NO.: 16), AHK3 (SEQ ID NO.: 17), or CRE1(AHK4/WOL) (SEQ ID NO.: 18) were cotransfected into *Arabidopsis* protoplast with the cytokinin-inducible reporter *ARR6-LUC*. Specifically, protoplasts were cotransfected with the *ARR6-LUC* reporter and the effector plasmid expressing AHK2, AHK3, CRE1(AHK4/WOL), CRE1(H467Q) (SEQ ID NO.: 19), CRE1(D973N) (SEQ ID NO.: 20), or mER7-GFP (control), and were analyzed after 6 hours of cytokinin treatment. Different from the CKI1 activation, over-expression of AHK2, AHK3, or CRE1(AHK4/WOL) did not enhance the expression of *ARR6-LUC* from the basal level in the absence of exogenous cytokinin. However, cytokinin treatment resulted in further activation of the reporter, especially for CRE1(AHK4/WOL) (Fig.1E). The requirement of the conserved His 467 and Asp 973 residues in the HK and the receiver domain, respectively, for the CRE1 activity was also tested. In Fig. 1E top, the expression of *Arabidopsis* HK proteins was detected by [³⁵S]methionine-labeled AHK2 (SEQ ID NO.: 26), AHK3 (SEQ ID NO.: 27), CRE1(AHK4/WOL) (SEQ ID NO.: 28), CRE1(H467Q) (SEQ ID NO.: 29), and CRE1(D973N) (SEQ ID NO.: 30).

30) that were immunoprecipitated and analyzed by SDS-PAGE. CRE1(H467Q) and CRE1(D973N) mutants lost their ability to further enhance *ARR6-LUC* expression in the presence of exogenous cytokinin (Fig. 1E).

Moreover, both mutants also imposed a dominant negative effect on blocking cytokinin signalling. Similar to the CKI1 mutants, the His mutant is more potent than the Asp mutant, perhaps due to the flexible phosphorelay process between mutant and wild-type HKs. These results indicated that cytokinin signals are sensed by multiple HK receptors with different or overlapping expression patterns. Although CRE1 is predominantly expressed in roots (Mahonen et al., *supra*; Ueguchi et al., *supra*), it can certainly function in leaf cells as a cytokinin receptor. However, CKI1 and CRE1 represent two different types of cytokinin receptors that require HK activity and phosphorelay to initiate cytokinin signalling, but have different regulatory mechanisms and probably distinct affinity or specificity for cytokinins.

To gain insight into where the cytokinin signalling is perceived, the sub-cellular localization of CKI1 was examined using the CKI1-GFP fusion. We first confirmed that CKI1-GFP acted similarly as CKI1-HA in the protoplast assay based on *ARR6-LUC* activation (Fig. 1D). Confocal microscopy showed that CKI1 is mainly localized to the plasma membrane but not in the cytosol, ER or nucleus, indicated by various GFP markers (Fig. 1F). Specifically, protoplasts were transfected with CKI1-GFP or various GFP marker plasmid DNA. Tunicamycin (1 µg/ml) treatment was for 12 hours. The transfected protoplasts were analyzed with a confocal microscope. The subcellular GFP markers are “CAT-GFP” for the cytosol, “ER-GFP” for the endoplasmic reticulum, and “N-GFP” (HSP-GFP) for the nucleus. Since the N-terminus of CKI1 is replete with putative glycosylation sites, the effect of a glycosylation inhibitor, tunicamycin, in CKI1-GFP expression, localization or function was examined. Although tunicamycin did not affect the expression of other GFP constructs, the expression

of CKI1-GFP was completely abolished (Fig. 1F). This result showed that the glycosylation of CKI1 is important for its stability, processing and/or trafficking in plant cells.

5 HPs Acts as a Shuttle Between the Cytosol and Nucleus Shuttles in Cytokinin Signalling

The studies of multistep phosphorelay in yeast osmosensing and in various prokaryotic systems have shown the use of histidine-containing phosphotransmitters (HP) as signal transducers between receptors and response regulators (Stock et al., *supra*; Posas et al., *supra*; Uhl and Miller, *EMBO J.* 15: 1028-36, 1996). The analysis of the *Arabidopsis* genome has revealed at least five genes encoding putative AHPs with unknown physiological functions. To determine whether AHPs are involved in the cytokinin signal transduction, AHP1, AHP2, and AHP5 were cloned into the plant expression vector and tested in the protoplast cytokinin response assay based on the *ARR6-LUC* reporter activity. AHP1, AHP2, and AHP5 slightly increased the expression of the *ARR6-LUC* with or without *t*-zeatin (Fig. 2A). Specifically, protoplasts were cotransfected with the *ARR6-LUC* reporter and an effector plasmid carrying AHP1 (SEQ ID NO.: 21), AHP2 (SEQ ID NO.: 22), AHP5 (SEQ ID NO.: 23), or AHP1 mutants (SEQ ID NOS.: 24 and 25). Transfected protoplasts were incubated for 6 hours to allow effector expression before cytokinin treatment for 3 hours. The AHP1 mutant, where the conserved His79 was replaced with Gln, did not repress the cytokinin activation of *ARR6-LUC*. Even the AHP1(H77Q, H79Q; SEQ ID NO.: 25) double mutant was not able to inhibit the *ARR6-LUC* activation by cytokinin. AHPs and AHP1 mutants were expressed at comparable levels in the protoplast system (Fig. 2A top). This result indicated that AHPs are not the limiting factor or the key switch/regulators in cytokinin signalling. Their action as potential mediators may require cytokinin-dependent modification.

Previous *in vitro* studies have suggested that AHP1 and AHP2 could interact with the receiver domain of CKI1 (Nakamura et al., *Biosci. Biotechnol. Biochem.* 63: 1627-30, 1999).

To gain further *in vivo* support for the involvement of AHPs in cytokinin signalling, the action of AHP1-GFP in transfected protoplast in the absence or presence of cytokinin was followed using fluorescence microscopy. Specifically, protoplasts were transfected with *AHP1-GFP*, *AHP2-GFP*, or *AHP5-GFP* and incubated for 3 hours to allow sufficient expression. After *t*-zeatin (100 nM) treatment for 30 minutes, transfected cells were observed under a fluorescence microscope. As shown in Fig. 2B, AHP1-GFP was mainly localized in the cytoplasm without cytokinin stimulation, but was translocated into the nucleus by *t*-zeatin treatment. The cytokinin-dependent translocation of AHP1-GFP was transient, occurring within 30 to 90 minutes after cytokinin treatment (Fig. 2B). AHP2-GFP showed similar cytokinin-dependent translocation but not AHP5-GFP, suggesting their different functions in plant cells. Thus, AHP1 and AHP2 are believed to serve as shuttles and phosphorelay carriers between the cytokinin receptors and the downstream nuclear responses.

Opposite Regulation of Early Cytokinin Response Gene Transcription by Two Types of Nuclear RRs in Cytokinin Signalling

The requirement of HK activity and phosphorelay in the initiation of cytokinin signalling from CKI1 and CRE1, the cytokinin-inducible *ARR6* expression, the cytokinin-dependent translocation of AHPs into the nucleus, and the physical interaction of AHPs and ARRs in yeast (Suzuki et al., *Plant Cell Physiol.* 42: 37-45, 2001) suggest that some ARRs may act downstream in cytokinin signal transduction. There are two major subfamilies of *RR* genes in the *Arabidopsis* genome: the cytokinin-inducible A type and the DNA-binding B type (Urao, et al., *supra*; D'Agostino and Kieber, *Trends Biochem. Sci.* 24: 452-6, 1999; Sakai et al., *Plant J.* 24: 703-11, 2000). To investigate whether any *Arabidopsis* RRs are involved in cytokinin signalling, four representative A-type

ARR genes and three B-type *ARR* genes were cloned into a standard plant expression vector with an HA or GFP tag. Specifically, protoplasts were cotransfected with the *ARR6-LUC* reporter and the effector plasmid carrying *ARR1*, *ARR2*, *ARR4*, *ARR5*, *ARR6*, *ARR7*, *ARR10*, or various mutants, *ARR2* (D80N) (SEQ ID NO.: 10), *ARR4*(D95N) (SEQ ID NO.: 11), and *ARR6*(D76N) (SEQ ID NO.: 12). As shown in Fig. 3A, A-type ARRs, such as *ARR4*, *ARR5*, *ARR6*, and *ARR7*, repressed *ARR6-LUC* activity induced by 100 nM *t*-zeatin. Although *ARR6* seemed to be expressed at lower abundance, its repression activity was the strongest.

10 In contrast, the B-type ARRs, such as *ARR1*, *ARR2*, and *ARR10*, dramatically activated *ARR6-LUC* expression. Over-expression of *ARR1*, *ARR2*, and *ARR10* was sufficient to activate cytokinin signalling at different levels in the absence of exogenous cytokinin. *ARR1* and *ARR2* activated *ARR6-LUC* about 40- and 400-fold, respectively. Cytokinin treatment further enhanced the effect of *ARR2* on *ARR6-LUC* to over 1000-fold. The lower activation by *ARR1* could be due to its lower expression level in transfected protoplasts. In the absence of cytokinin, *ARR10* could activate *ARR6-LUC* about 10 fold. Cytokinin enhanced the effect of *ARR10* on the *ARR6-LUC* activity another 10-20 fold. The differential effect of *ARR2* and *ARR10* in activating cytokinin signalling could be attributed to differences in their intrinsic activities in DNA binding and/or transcription activation. It could also be due to their distinct affinity to endogenous repressors that likely exist in leaf protoplasts to prevent cytokinin signalling under unstimulated condition. Surprisingly, mutations of the conserved Asp residue in the receiver domains of *ARR4*(D95N), *ARR6*(D76N), and *ARR2*(D80N) did not lead to alteration of their repression and activation functions, respectively, in cytokinin signalling (Fig. 3A). The transfected cells were incubated for 3 hours to allow effector protein expression. This result is consistent with the model that phosphorelay stimulated by cytokinin may be involved in de-repression of the positive regulators like *ARR1*, *ARR2*, and

ARR10. Over-expression of these ARR2s likely bypasses the negative regulation and causes constitutive cytokinin responses without the signal.

To further support the role of ARR2 in cytokinin signalling, we designed a dominant mutant of ARR2 (Δ ARR2) (SEQ ID NO.: 4) containing only the DNA
5 binding domain but not the putative transcription activation and the receiver domains (Sakai et al., *supra*). If Δ ARR2 can compete with the endogenous ARR2s in binding to the *ARR6* promoter without transcription activation, it may block cytokinin- or CKI1-dependent activation of *ARR6-LUC*. In Fig. 3B, protoplasts were cotransfected with the *ARR6-LUC* reporter alone or with the effector
10 plasmids as indicated. Transfected protoplasts were incubated for 6 hours to allow effector protein expression before treatment with 100 nM *t*-zeatin for 3 hours. As shown in Fig. 3B, the dominant negative mutant of ARR2 could effectively block *ARR6-LUC* expression in the presence of 100 nM *t*-zeatin or CKI1, suggesting that ARR2 and/or ARR2-like proteins act downstream of
15 cytokinin and CKI1-initiated phosphorelay as the nuclear targets in the cytokinin signal transduction pathway. Furthermore, the dominant negative effect of the CKI1(H450Q) mutant on the protoplast response to exogenous cytokinin could be bypassed by over-expression of the wild-type ARR2 (Fig. 3B). These epistasis analyses place ARR2 downstream of multiple cytokinin HK receptors. The
20 finding of multiple ARR2 binding motifs, (G/A)GAT(T/C), in the promoter regions of *ARR6* and other cytokinin-inducible genes suggests that ARR2 could be a master regulator in cytokinin signalling (*Id.*). Exogenous cytokinin-initiated phosphorelay through CRE1, or AHK2 and AHK3 likely converges with the CKI1 phosphorelay on the common AHP shuttles and the nuclear ARR
25 transcription activators.

To provide more evidence that the A-type and B-type ARR2s are transcription repressors and activators, respectively, in cytokinin signalling, the sub-cellular localization of ARR6-GFP and ARR2-GFP was examined. In Fig. 3C, Protoplasts were transfected with *ARR2-GFP*, *ARR2(D80N)-GFP*, Δ *ARR2*-
30 *GFP*, or *ARR6-GFP* plasmid DNA and observed after 3-6 hours of incubation

with a fluorescence microscope. Both types of ARRs are exclusively localized in the nucleus regardless of the cytokinin treatment (Fig. 3C). Their nuclear localization is likely independent of the phosphorylation state because ARR2(D80N)-GFP was found in the nucleus. The ARR2 dominant negative mutant (Δ ARR2) was also localized in the nucleus (Fig. 3C). Thus, the cytokinin-dependent phosphorelay does not play a role in the nuclear localization and DNA-binding of ARRs or their intrinsic transcription activation or repression activities. The results support the model that cytokinin-initiated phosphorelay plays a role in releasing the sequestered activation-type ARR transcription factors from the yet unknown repressors (Fig. 5).

As depicted in Fig. 5, cytokinin signal is perceived internally or externally by multiple HKs at the plasma membrane (AHK2, AHK3, CKI1, or CRE1). Upon perception of the cytokinin signal, HKs initiate a signalling cascade via the phosphorelay that in turn results in the nuclear translocation of AHPs from the cytosol. Activated AHPs interact with sequestered ARRs or ARR complex and release the activation type of ARRs from putative repressor in the nucleus. The de-repressed ARRs bind to multiple *cis* elements in the promoter of target genes. The transcription activation is essential for cell proliferation, shoot formation, and delayed senescence. The activation of the repressor-type of ARRs as early cytokinin response genes provides a negative feedback mechanism that allows plant cells for fine-tuning or resetting the physiological state in diverse differentiation processes during plant growth and development (RD, response domain; BD, DNA binding domain; AD, transcription activation domain; PM, plasma membrane; N, nucleus; R, repressor).

This model is consistent with the findings that the N-terminus of ARR1 or ARR2 inhibits transcription based on a synthetic promoter (*Id.*), and the addition of cytokinin further enhances the effects of ARR1, ARR2 and ARR10 over-expression on the *ARR6-LUC* activity (Fig. 3). The activation of the repressor-

type of ARRs is thought to provide a negative feedback loop in controlling the transient induction of cytokinin early response genes and allows resetting and/or fine-tuning of the physiological state of the cells.

5 Ectopic expression of the master regulator ARR2 mimics cytokinin responses

In tissue culture, induction of cell proliferation and subsequent shoot formation require cytokinin. To determine whether the same cytokinin signalling pathway is responsible for transcription regulation, cell proliferation, shoot meristem initiation, and leaf formation, we developed a seedling cytokinin
10 response assay. In Figs. 4A and 4B, *Arabidopsis* seedlings were transformed with *Agrobacteria* carrying *GFP*, *CKI1*, *CKI1(H405Q)*, *ARR2*, and *ARR6* constructs in the mini-binary vector pCB302 (Xiang et al., *Plant Mol. Biol.* 40: 711-7, 1999). Seedlings transformed with the *Agrobacteria* carrying the binary vector pCB301 without a Bar selection marker were used as a transformation control (vector).
15 The transformed seedlings were selected with glufosinate ammonium (10 µg/ml) and maintained on medium with auxin (IAA) only (Fig. 4A, +IAA, -2IP) or with auxin and cytokinin (Fig. 4B, +IAA, +2IP) for 14 days. The transformed tissues were mostly derived from the shoot apical meristem. The GFP control showed green callus formation with exogenous IAA but promoted shoot formation with
20 exogenous cytokinin. Both CKI1 and ARR2 promoted extensive cell proliferation and shoot formation even without exogenous cytokinin. The experiments were repeated three times with similar results. Compared with the GFP control, the CKI1(H405Q) mutant and ARR6 slightly inhibited cell proliferation on the same medium without or with exogenous cytokinin (Figs. 4A
25 and 4B). The results of this cytokinin seedling assay are consistent with those from protoplast transient expression analyses based on transcription. It is therefore clear that the HK activity of CKI1 is required to perceive and/or initiate cytokinin signalling.

To observe the consequences of constitutive cytokinin signalling *in planta*,
30 the binary vectors expressing CKI1, CKI1(H405Q), ARR2, and ARR6 were

introduced into *Arabidopsis* plants. Ectopic expression of ARR2 in transgenic *Arabidopsis* plants displayed a spectrum of cytokinin-related phenotypes, including reduced apical dominance, increased leaf number and size, altered leaf shape and color, and delayed leaf senescence (Fig. 4C). Transgenic *Arabidopsis* plants carrying the CKI1 construct showed similar phenotypes, but not the CKI1(H405Q) and ARR6 transgenic plants serving as controls. Taken together, CKI1 and ARR2 are positive regulators in the cytokinin signal transduction pathway mediating transcription, cell proliferation, and shoot and leaf formation, as well as delaying leaf senescence. ARR2 could be one of the direct downstream targets of CKI1 and CRE1, and serve as a master regulator in diverse cytokinin responses.

Materials and Methods

The above-mentioned results were obtained using the following materials and methods.

Plasmid Constructs

The 2.4 kb *Arabidopsis* ARR6 promoter was fused to the firefly luciferase gene to create the ARR6-LUC reporter construct. The CKI1 and AHK2 genes were amplified from the *Arabidopsis* genomic DNA by polymerase chain reaction (PCR). The AHK3, CRE1(CRE1/WOL), AHP1, AHP2, AHP5 and ARR1, ARR 2, ARR4, ARR5, ARR6, ARR7, and ARR10 coding regions were obtained by PCR from an *Arabidopsis* cDNA library (Sheen, *Science* 274: 1900-2, 1996). All of the mutants were generated by QuickChange site-directed mutagenesis (Stratagene). The coding regions of all proteins were tagged with either two copies of the haemagglutinin epitope (DHA) or GFP and inserted into a plant expression vector containing the 35S4PPDK promoter (Sheen, *supra*) and the nos terminator. All PCR products and the mutations were confirmed by DNA sequencing.

Arabidopsis Protoplast Transient Expression Assay

Arabidopsis Bensheim (BE) protoplasts were isolated and transfected as described with some modifications (Kovtun et al., *Proc. Natl. Acad. Sci. U S A* 97: 2940-5 2000). Typically, 2×10^4 protoplasts were transfected with 10 μ g of
5 plasmid carrying different combinations of a reporter, effectors, and an internal control. For instance, in the experiments shown in Fig. 1A, protoplasts (4×10^4) were transfected with 2 μ g of *UBQ10-GUS* (internal control) and 20 μ g of *ARR6-LUC*, *GH3-LUC*, or *RD29A-LUC* plasmid DNA. Transfected protoplasts were incubated at 1×10^4 per ml with/without 1-100 nM of *t*-zeatin for 3-6 hours under
10 the light at 23 °C. The *UBQ10-GUS* construct was used as an internal control to normalize the variations of each transfection in cell numbers, transformation efficiency and cell viability. The results are shown as the means of duplicate samples with the standard deviation. All transient experiments were repeated at least three times with similar results. GFP fluorescence was observed by either
15 Nikon TE200 fluorescent microscopy or Leica TCSNT confocal microscopy.

Protein Immunoprecipitation

Transfected protoplasts were incubated with [35 S]-methionine (200 μ Ci/ml) for 6 hours. The effector proteins were immunoprecipitated as described (Sheen,
20 *supra*), analyzed by SDS/PAGE (10 or 12.5 %) and visualized by fluorography.

Seedling Cytokinin Response Assay

Four-day-old *Arabidopsis* BE seedlings were placed on the callus induction medium, containing 0.5 mg/L 2,4-D, 0.05 mg/L BAP, and 0.05 mg/L
25 kinetin, for four days. The seedlings were co-cultivated and transformed with *Agrobacterium tumefaciens* GV3101 carrying pCB302 with either *CKI1*, *CKI1(H405Q)*, *ARR2*, *ARR6*, *GFP* under control of the 35SC4PPDK promoter, or the pCB301 binary vector (Xiang et al., *Plant Mol. Biol.* 40:711-7, 1999) alone for three days. Transformed seedlings were then selected on the glufosinate
30 ammonium-containing medium (with either 0.15 mg/L IAA alone or with 0.15

mg/L IAA and 5 mg/L 2IP) and observed up to 4 weeks. The transformed proliferating tissues were mostly derived from the apical shoot meristem of the seedlings.

5 Transgenic plant analysis

The same constructs analyzed in the *Arabidopsis* protoplast transient expression and seedling assays, including *CKI1*, *CKI1(H405Q)*, *ARR2*, and *ARR6*, were used to generate *Arabidopsis* transgenic plants using the floral dip method and Bar selection as described (Clough and Bent, *Plant J.* 16: 735-43, 1998).

- 10 More than 200 transgenic plants were obtained with each construct. The *ARR2* and *CKI1* transgenic plants developed branches early, produced more leaves that are bigger and greener but flowered later than the control *CKI1(H405Q)* and *ARR6* transgenic plants. For senescence assay, fully expanded fourth leaves from representative transgenic plants were detached and floated on distilled water in
15 the dark for four days.

Isolation of Sequences Encoding A-type and B-type Response Regulators or Histidine Kinases

- The isolation of additional genes encoding A-type and B-type response
20 regulators or histidine kinase genes is accomplished using standard strategies and techniques that are well known in the art.

- In one particular example, the B-type response regulator sequence (for example, *ARR1*, *ARR2*, or *ARR10* sequences), the A-type response regulator sequence, (for example, *ARR4*, *ARR5*, *ARR6*, or *ARR7*), or the histidine kinase
25 (for example, *CRE1* or *CKI1*), described herein may be used, together with conventional screening methods of nucleic acid hybridization screening, to isolate additional sequences encoding these regulators. Such hybridization techniques and screening procedures are well known to those skilled in the art and are described, for example, in Benton and Davis, *Science* 196: 180, 1977; Grunstein
30 and Hogness, *Proc. Natl. Acad. Sci., USA* 72: 3961, 1975; Ausubel et al. *Current*

Protocols in Molecular Biology, Wiley Interscience, New York; Berger and Kimmel, *Guide to Molecular Cloning Techniques*, 1987, Academic Press, New York.; and Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, New York. In one particular example, all or part
5 of the response regulator gene may be used as a probe to screen a recombinant plant DNA library for genes having sequence identity or similarity to the response regulator gene. Hybridizing sequences are detected by plaque or colony hybridization according to the methods described below.

Alternatively, using all or a portion of the amino acid sequence of a
10 response regulator, one may readily design oligonucleotide probes, including degenerate oligonucleotide probes (i.e., a mixture of all possible coding sequences for a given amino acid sequence). These oligonucleotides may be based upon the sequence of either DNA strand and any appropriate portion of the response regulator sequence. General methods for designing and preparing such
15 probes are provided, for example, in Ausubel et al., *supra*; and Berger and Kimmel, *supra*. These oligonucleotides are useful for response regulator or histidine kinase sequence isolation, either through their use as probes capable of hybridizing to response regulator complementary sequences or as primers for various amplification techniques, for example, polymerase chain reaction (PCR)
20 cloning strategies. If desired, a combination of different oligonucleotide probes may be used for the screening of a recombinant DNA library. The oligonucleotides may be detectably-labeled using methods known in the art and used to probe filter replicas from a recombinant DNA library. Recombinant DNA libraries are prepared according to methods well known in the art, for
25 example, as described in Ausubel et al. (*supra*), or they may be obtained from commercial sources.

As discussed above, response regulator-specific oligonucleotides may also be used as primers in amplification cloning strategies, for example, using PCR. PCR methods are well known in the art and are described, for example, in *PCR*
30 *Technology*, Erlich, ed., Stockton Press, London, 1989; *PCR Protocols: A Guide*

to *Methods and Applications*, Innis et al., eds., Academic Press, Inc., New York, 1990; and Ausubel et al. (*supra*). Primers are optionally designed to allow cloning of the amplified product into a suitable vector, for example, by including appropriate restriction sites at the 5' and 3' ends of the amplified fragment (as
5 described herein). If desired, response regulator sequences may be isolated using the PCR "RACE" technique, or Rapid Amplification of cDNA Ends (see, e.g., Innis et al. (*supra*)). By this method, oligonucleotide primers based on the response regulator sequence are oriented in the 3' and 5' directions and are used to generate overlapping PCR fragments. These overlapping 3'- and 5'-end RACE
10 products are combined to produce an intact full-length cDNA. This method is described in Innis et al. (*supra*); and Frohman et al., *Proc. Natl. Acad. Sci. USA* 85: 8998, 1988.

Confirmation of a sequence's relatedness to the response regulators may be accomplished by a variety of conventional methods including, but not limited to,
15 sequence comparison of the gene and its expressed product to known response regulators for example those disclosed herein. In addition, the activity of the gene product may be evaluated according to any of the techniques described herein.

Once a response regulator gene is identified, it is cloned according to
20 standard methods and used for the construction of plant expression vectors as described below.

Expression Constructs

Those skilled in the field of molecular biology will understand that any
25 of a wide variety of expression systems may be used to express the response regulators disclosed herein. For example, an A-type or B-type response regulator may be expressed for the desired effect in any of a number of plant hosts including, without limitation, algae, tree species, ornamental species, temperate fruit species, tropical fruit species, vegetable species, legume species, crucifer
30 species, monocots, dicots, or in any plant of commercial or agricultural

significance. Particular examples of suitable plant hosts include, but are not limited to, Conifers, Petunia, Tomato, Potato, Tobacco, *Arabidopsis*, Lettuce, Sunflower, Oilseed rape, Flax, Cotton, Sugarbeet, Celery, Soybean, Alfalfa, *Medicago*, Lotus, *Vigna*, Cucumber, Carrot, Eggplant, Cauliflower, Horseradish, Morning Glory, Poplar, Walnut, Apple, Grape, Asparagus, Rice, Maize, Millet, Onion, Barley, Orchard grass, Oat, Rye, and Wheat. In addition, as is discussed below, transgenic expression constructs may be expressed in a plant to increase plants yield, shoot formation, leaf growth, photosynthesis, or delay senescence.

Materials for expressing these genes are available from a wide range of sources including the American Type Culture Collection (Rockland, MD); or from any of a number seed companies, for example, W. Atlee Burpee Seed Co. (Warminster, PA), Park Seed Co. (Greenwood, SC), Johnny Seed Co. (Albion, ME), or Northrup King Seeds (Harstville, SC). Descriptions and sources of useful host cells are also found in Vasil, Cell Culture and Somatic Cell Genetics of Plants, Vol I, II, III Laboratory Procedures and Their Applications Academic Press, New York, 1984; Dixon, Plant Cell Culture-A Practical Approach, IRL Press, Oxford University, 1985; Green et al., Plant Tissue and Cell Culture, Academic Press, New York, 1987; and Gasser and Fraley, Science 244: 1293, 1989.

The method of transformation or transfection and the choice of vehicle for expression of the A-type or B-type response regulators, or a histidine kinase will depend on the host system selected. Transformation and transfection methods are described, e.g., in Ausubel et al. (*supra*); Weissbach and Weissbach, Methods for Plant Molecular Biology, Academic Press, 1989; Gelvin et al., Plant Molecular Biology Manual, Kluwer Academic Publishers, 1990; Kindle, *Proc. Natl. Acad. Sci., USA* 87: 1228, 1990; Potrykus, *Annu. Rev. Plant Physiol. Plant Mol. Biology* 42: 205, 1991; and BioRad (Hercules, CA) Technical Bulletin #1687 (Biolistic Particle Delivery Systems). Expression vehicles may be chosen from those provided, e.g., in Cloning Vectors: A Laboratory Manual (Pouwels et al., 1985, Supp. 1987); Gasser and Fraley (*supra*); Clontech Molecular Biology

Catalog (Catalog 1992/93 Tools for the Molecular Biologist, Palo Alto, CA); and the references cited above. Other expression constructs are described by Fraley et al. (U.S. Pat. No. 5,352,605).

Most preferably, an A-type or B-type response regulator or a histidine
5 kinase is produced by a stably-transfected plant cell line, a transiently-transfected plant cell line, or by a transgenic plant. A number of vectors suitable for stable transfection of plant cells or for the establishment of transgenic plants are available to the public; such vectors are described in Pouwels et al. (*supra*), Weissbach and Weissbach (*supra*), and Gelvin et al. (*supra*). Methods for
10 constructing such cell lines are described in, e.g., Weissbach and Weissbach (*supra*), and Gelvin et al. (*supra*). Typically, plant expression vectors include (1) a cloned plant gene under the transcriptional control of 5' and 3' regulatory sequences and (2) a dominant selectable marker. Such plant expression vectors may also contain, if desired, a promoter regulatory region (for example, one
15 conferring inducible or constitutive, pathogen- or wound-induced, environmentally- or developmentally-regulated, or cell- or tissue-specific expression), a transcription initiation start site, a ribosome binding site, an RNA processing signal, a transcription termination site, and/or a polyadenylation signal.

20 Once the desired nucleic acid sequence encoding an A-type or B-type response regulators or a histidine kinase is obtained as described above, it may be manipulated in a variety of ways known in the art. For example, where the sequence involves non-coding flanking regions, the flanking regions may be subjected to mutagenesis.

25 In general, the constructs will involve regulatory regions functional in plants which provide for modified production of the regulator protein or regulator transcript (such as an antisense transcript) as discussed herein. The open reading frame coding for the regulator protein or functional fragment thereof will be joined at its 5' end to a transcription initiation regulatory region such as the
30

sequence naturally found in the 5' upstream region of the A-type or B-type response regulator, or a histidine kinase. Numerous other transcription initiation regions are available which provide for constitutive or inducible regulation.

For applications where developmental, cell, tissue, hormonal, or environmental expression is desired, appropriate 5' upstream non-coding regions are obtained from other genes, for example, from genes regulated during meristem development, seed development, embryo development, or leaf development.

Regulatory transcript termination regions may also be provided in DNA constructs of this invention as well. Transcript termination regions may be provided by the DNA sequence encoding the A-type or B-type response regulator or histidine kinase, or any convenient transcription termination region derived from a different gene source. The transcript termination region will contain preferably at least 1-3 kb of sequence 3' to the structural gene from which the termination region is derived. Plant expression constructs having, for example, A-type or B-type response regulator as the DNA sequence of interest for expression may be employed with a wide variety of plant life. Such genetically-engineered plants are useful for a variety of industrial and agricultural applications as discussed herein. Importantly, this invention is applicable to dicotyledons and monocotyledons, and will be readily applicable to any new or improved transformation or regeneration method.

An example of a useful plant promoter according to the invention is a caulimovirus promoter, for example, a cauliflower mosaic virus (CaMV) promoter. These promoters confer high levels of expression in most plant tissues, and the activity of these promoters is not dependent on virally encoded proteins. CaMV is a source for both the 35S and 19S promoters. In most tissues of transgenic plants, the CaMV 35S promoter is a strong promoter (see, e.g., Odell et al., *Nature* 313: 810, 1985). The CaMV promoter is also highly active in monocots (see, e.g., Dekeyser et al., *Plant Cell* 2: 591, 1990; Terada and Shimamoto, *Mol. Gen. Genet.* 220: 389, 1990). Moreover, activity of this

promoter can be further increased (i.e., between 2-10 fold) by duplication of the CaMV 35S promoter (see e.g., Kay et al., *Science* 236: 1299, 1987; Ow et al., *Proc. Natl. Acad. Sci., U.S.A.* 84: 4870, 1987; and Fang et al., *Plant Cell* 1: 141, 1989). In addition, the a minimal 35S promoter may also be used as is described
5 herein.

Other useful plant promoters include, without limitation, the nopaline synthase promoter (An et al., *Plant Physiol.* 88: 547, 1988) and the octopine synthase promoter (Fromm et al., *Plant Cell* 1: 977, 1989).

For certain applications, it may be desirable to produce the regulator in
10 an appropriate tissue, at an appropriate level, or at an appropriate developmental time. For this purpose, there are an assortment of gene promoters, each with its own distinct characteristics embodied in its regulatory sequences, shown to be regulated in response to the environment, hormones, and/or developmental cues. These include gene promoters that are responsible for heat-regulated gene
15 expression (see, e.g., Callis et al., *Plant Physiol.* 88: 965, 1988; Takahashi and Komeda, *Mol. Gen. Genet.* 219: 365, 1989; and Takahashi et al., *Plant J.* 2: 751, 1992), light-regulated gene expression (e.g., the *Arabidopsis Cab2* photosynthetic, leaf specific promoter described by Mitra et al., *Plant Mol. Biol.* 12: 169-179, 1989; the pea *rbcS-3A* described by Kuhlemeier et al., *Plant Cell* 1:
20 471, 1989; the maize *rbcS* promoter described by Schöffner and Sheen, *Plant Cell* 3: 997, 1991; or the cholorophyll a/b-binding protein gene found in pea described by Simpson et al., *EMBO J.* 4: 2723, 1985), hormone-regulated gene expression (for example, the abscisic acid (ABA) responsive sequences from the *Em* gene of wheat described by Marcotte et al., *Plant Cell* 1: 969, 1989; the ABA-inducible
25 HVA1 and HVA22, and rd29A promoters described for barley and *Arabidopsis* by Straub et al., *Plant Cell* 6: 617, 1994, Shen et al., *Plant Cell* 7: 295, 1995), wound-induced gene expression (for example, of *wun1* described by Siebertz et al., *Plant Cell* 1: 961, 1989), and organ-specific gene expression (for example, of the tuber-specific storage protein gene described by Roshal et al., *EMBO J.* 6:
30 1155, 1987; the 23-kDa zein gene from maize described by Schernthaner et al.,

EMBO J. 7: 1249, 1988; or the French bean β -phaseolin gene described by Bustos et al., *Plant Cell* 1: 839, 1989; the vegetative storage protein promoter (soybean vspB) described by Sadka et al. (*Plant Cell* 6: 737-749, 1994)), cycling promoters (e.g., the *Arabidopsis* cdc2a promoter described by Hemerly et al., *Proc. Natl. Acad. Sci. USA* 89: 3295-3299, 1992), senescence-specific promoters (e.g., the *Arabidopsis* SAG12 promoter described by Gan et al, *Science*: 270, 1986-1988, 1995), seed-specific promoters (for example, endosperm-specific or embryo-specific promoters), or pathogen-inducible promoters (for example, PR-1 or β -1,3 glucanase promoters).

10 Plant expression vectors may also optionally include RNA processing signals, e.g, introns, which have been shown to be important for efficient RNA synthesis and accumulation (Callis et al., *Genes and Dev.* 1: 1183, 1987). The location of the RNA splice sequences can dramatically influence the level of transgene expression in plants. In view of this fact, an intron may be positioned
15 upstream or downstream of the regulator encoding sequence in the transgene to modulate levels of gene expression.

In addition to the aforementioned 5' regulatory control sequences, the expression vectors may also include regulatory control regions which are generally present in the 3' regions of plant genes (Thornburg et al., *Proc. Natl. Acad. Sci. U.S.A.* 84: 744, 1987; An et al., *Plant Cell* 1: 115, 1989). For
20 example, the 3' terminator region may be included in the expression vector to increase stability of the mRNA. One such terminator region may be derived from the PI-II terminator region of potato. In addition, other commonly used terminators are derived from the octopine or nopaline synthase signals.

25 The plant expression vector also typically contains a dominant selectable marker gene used to identify those cells that have become transformed. Useful selectable genes for plant systems include genes encoding antibiotic resistance, for example, those encoding resistance to hygromycin, kanamycin, bleomycin, G418, streptomycin, or spectinomycin. Genes required for
30 photosynthesis may also be used as selectable markers in photosynthetic-deficient

strains. Finally, genes encoding herbicide resistance may be used as selectable markers; useful herbicide resistance genes include the *bar* gene encoding the enzyme phosphinothricin acetyltransferase and conferring resistance to the broad spectrum herbicide Basta® (Hoechst AG, Frankfurt, Germany).

5 Efficient use of selectable markers is facilitated by a determination of the susceptibility of a plant cell to a particular selectable agent and a determination of the concentration of this agent which effectively kills most, if not all, of the untransformed cells. Some useful concentrations of antibiotics for tobacco transformation include, e.g., 75-100 µg/mL (kanamycin), 20-50 µg/mL
10 (hygromycin), or 5-10 µg/mL (bleomycin). A useful strategy for selection of transformants for herbicide resistance is described, e.g., by Vasil et al., *supra*.

 It should be readily apparent to one skilled in the art of molecular biology, especially in the field of plant molecular biology, that the level of gene expression is dependent, not only on the combination of promoters, RNA
15 processing signals, and terminator elements, but also on how these elements are used to increase the levels of selectable marker gene expression.

Plant Transformation

 Upon construction of the plant expression vector, several standard
20 methods are available for introduction of the vector into a plant host, thereby generating a transgenic plant. These methods include (1) *Agrobacterium*-mediated transformation (*A. tumefaciens* or *A. rhizogenes*) (see, e.g., Lichtenstein and Fuller, In: *Genetic Engineering*, Vol 6, Rigby, ed., London, Academic Press, 1987; and Lichtenstein and Draper, In: *DNA Cloning*, Vol II, Glover, ed., Oxford,
25 IRI Press, 1985), (2) the particle delivery system (see, e.g., Gordon-Kamm et al., *Plant Cell* 2: 603, 1990); or BioRad Technical Bulletin #1687, *supra*), (3) microinjection protocols (see, e.g., Green et al., *supra*), (4) polyethylene glycol (PEG) procedures (see, e.g., Draper et al., *Plant Cell Physiol.* 23: 451, 1982; or e.g., Zhang and Wu, *Theor. Appl. Genet.* 76: 835, 1988), (5) liposome-mediated
30 DNA uptake (see, e.g., Freeman et al., *Plant Cell Physiol.* 25: 1353, 1984), (6)

electroporation protocols (see, e.g., Gelvin et al., *supra*; Dekeyser et al., *supra*; Fromm et al., *Nature* 319: 791, 1986; Sheen, *Plant Cell* 2: 1027, 1990; or Jang and Sheen, *Plant Cell* 6: 1665, 1994), (7) the vortexing method (see, e.g., Kindle, *supra*), and floral dip method (see, e.g., Clough and Bent, *Plant J.* 16, 735-43. (1998)). The method of transformation is not critical to the invention. Any method which provides for efficient transformation may be employed. As newer methods are available to transform crops or other host cells, they may be directly applied.

The following is an example outlining one particular technique, an *Agrobacterium*-mediated plant transformation. By this technique, the general process for manipulating genes to be transferred into the genome of plant cells is carried out in two phases. First, cloning and DNA modification steps are carried out in *E. coli*, and the plasmid containing the gene construct of interest is transferred by conjugation or electroporation into *Agrobacterium*. Second, the resulting *Agrobacterium* strain is used to transform plant cells. Thus, for the generalized plant expression vector, the plasmid contains an origin of replication that allows it to replicate in *Agrobacterium* and a high copy number origin of replication functional in *E. coli*. This permits facile production and testing of transgenes in *E. coli* prior to transfer to *Agrobacterium* for subsequent introduction into plants. Resistance genes can be carried on the vector, one for selection in bacteria, for example, streptomycin, and another that will function in plants, for example, a gene encoding kanamycin resistance or herbicide resistance. Also present on the vector are restriction endonuclease sites for the addition of one or more transgenes and directional T-DNA border sequences which, when recognized by the transfer functions of *Agrobacterium*, delimit the DNA region that will be transferred to the plant.

In another example, plant cells may be transformed by shooting into the cell tungsten microprojectiles on which cloned DNA is precipitated. In the Biolistic Apparatus (Bio-Rad) used for the shooting, a gunpowder charge (22 caliber Power Piston Tool Charge) or an air-driven blast drives a plastic

macroprojectile through a gun barrel. An aliquot of a suspension of tungsten particles on which DNA has been precipitated is placed on the front of the plastic macroprojectile. The latter is fired at an acrylic stopping plate that has a hole through it that is too small for the macroprojectile to pass through. As a result, the plastic macroprojectile smashes against the stopping plate, and the tungsten microprojectiles continue toward their target through the hole in the plate. For the instant invention the target can be any plant cell, tissue, seed, or embryo. The DNA introduced into the cell on the microprojectiles becomes integrated into either the nucleus or the chloroplast.

In general, transfer and expression of transgenes in plant cells are now routine practices to those skilled in the art, and have become major tools to carry out gene expression studies in plants and to produce improved plant varieties of agricultural or commercial interest.

15 Transgenic Plant Regeneration

Whole plants can be regenerated, for example, from single cells, callus tissue, or leaf discs transformed with a plant expression vector according to standard plant tissue culture techniques. It is well known in the art that various cells, tissues, and organs from almost any plant can be successfully cultured to regenerate an entire plant; such techniques are described, e.g., in Vasil, *supra*; Green et al., *supra*; Weissbach and Weissbach, *supra*; and Gelvin et al., *supra*.

In one particular example, a cloned B-type response regulator or histidine kinase construct under the control of the *nos* promoter and the nopaline synthase terminator and carrying a selectable marker (for example, kanamycin resistance) is transformed into *Agrobacterium*. Transformation of leaf discs (for example, of tobacco or potato leaf discs), with vector-containing *Agrobacterium* is carried out as described by Horsch et al. (*Science* 227: 1229, 1985). Putative transformants are selected after a few weeks (for example, 3 to 5 weeks) on plant tissue culture media containing kanamycin (e.g. 100 µg/mL). Kanamycin-resistant shoots are then placed on plant tissue culture media without hormones

for root initiation. Kanamycin-resistant plants are then selected for greenhouse growth. If desired, seeds from self-fertilized transgenic plants can then be sowed in a soil-less medium and grown in a greenhouse. Kanamycin-resistant progeny are selected by sowing surfaced sterilized seeds on hormone-free kanamycin-
5 containing media. Analysis for the integration of the transgene is accomplished by standard techniques (see, for example, Ausubel et al., *supra*; Gelvin et al., *supra*).

Transgenic plants expressing the selectable marker are then screened for transmission of the transgene DNA by standard immunoblot and DNA
10 detection techniques. Each positive transgenic plant and its transgenic progeny are unique in comparison to other transgenic plants established with the same transgene. Integration of the transgene DNA into the plant genomic DNA is in most cases random, and the site of integration can profoundly affect the levels and the tissue and developmental patterns of transgene expression. Consequently,
15 a number of transgenic lines are usually screened for each transgene to identify and select plants with the most appropriate expression profiles.

Transgenic lines are evaluated for levels of transgene expression. Expression at the RNA level is determined initially to identify and quantitate expression-positive plants. Standard techniques for RNA analysis are employed
20 and include PCR amplification assays using oligonucleotide primers designed to amplify only transgene RNA templates and solution hybridization assays using transgene-specific probes (see, e.g., Ausubel et al., *supra*). The RNA-positive plants are then analyzed for protein expression by Western immunoblot analysis using specific antibodies (see, e.g., Ausubel et al., *supra*). In addition, *in situ*
25 hybridization and immunocytochemistry according to standard protocols can be done using transgene-specific nucleotide probes and antibodies, respectively, to localize sites of expression within transgenic tissue.

Increasing Plant Yield

To test the ability of the genes described herein to improve crop yield or productivity, seeds of transgenic plants expressing a B-type response regulator or histidine kinase are planted in test plots, and their agronomic performance is compared to standard plants using techniques familiar to those of skill in the art. 5 Optionally included in this comparison are plants of similar genetic background without the transgene. A yield benefit is observed and plants exhibiting the increased yield are advanced for commercialization.

In addition, transgenic plants expressing a B-type response regulator or histidine kinase are field tested for agronomic performance under conditions, 10 including, but not limited to, increased leaf production. When compared to nontransgenic plants, transgenic plants expressing the B-type response regulator exhibit higher yield than their non-transgenic counterparts.

15 Increasing Plant Shoot Formation

To test the ability of the genes described herein to increase shoot formation, seedlings are transformed with a transgene expressing a B-type response regulator or histidine kinase and shoot formation is measured, for example, by evaluating the development of green callus formation, using 20 techniques familiar to those of skill in the art. Optionally included in this comparison are plants of similar genetic background without the transgene. Plants exhibiting increased shoot formation relative to control plants are useful in the invention.

25 Delaying Plant Senescence

To test the ability of the genes described herein to delay senescence, fully expanded fourth leaves of plants expressing a recombinant B-type response regulator or histidine kinase are detached and floated on distilled water in the dark for four days, and their leaf size and quality of green color are compared

using techniques familiar to those of skill in the art. Optionally included in this comparison are plants of similar genetic background without the transgene.

Enhancing Plant Photosynthesis

5 To test the ability of the genes and domains described herein to enhance photosynthesis, seedlings are transformed with a transgene expressing a recombinant B-type response regulator or histidine kinase and photosynthesis measured using techniques familiar to those of skill in the art. Optionally included in this comparison are plants of similar genetic background without the
10 transgene.

Increasing Plant Leaf Growth

 To test the ability of the genes described herein to increase leaf growth, seedlings are transformed with a transgene expressing a recombinant B-type
15 response regulator and leaf growth measured, for example, by measuring leaf number and leaf area using techniques familiar to those of skill in the art. Optionally included in this comparison are plants of similar genetic background without the transgene.

20 Silencing A-type Response Regulator Gene Expression

 Plants having decreased expression of an A-type response regulator are useful, for example, for increasing plant productivity, delaying senescence, increasing photosynthesis, increased leaf growth, and increased productivity. Plants having decreased expression of such A-type response regulators are
25 generated according to standard gene silencing methods including, without limitation, co-suppression and antisense methodologies, expression of dominant negative gene products, and creation of plants having mutated genes encoding such regulators.

Co-Suppression

One preferred method of silencing gene expression is co-suppression (also referred to as sense suppression). This technique, which involves introduction of a nucleic acid configured in the sense orientation, has been shown to effectively
5 block the transcription of target genes (see for example, Napoli et al., *Plant Cell*, 2:279-289, 1990 and Jorgensen et al., U.S. Patent No. 5,034,323).

Generally, sense suppression involves transcription of the introduced sequence such as a gene encoding an A-type response regulator. However, co-suppression may also occur where the introduced sequence contains no coding
10 sequence per se, but only intron or untranslated sequences homologous to sequences present in the primary transcript of the endogenous gene to be repressed. The introduced sequence generally will be substantially identical to the endogenous gene targeted for repression. Such identity is typically greater than about 50%, but higher identities (for example, 80% or even 95%) are
15 preferred because they result in more effective repression. The effect of co-suppression may also be applied to other proteins within a similar family of genes exhibiting homology or substantial homology. Segments from a gene from one plant can be used directly, for example, to inhibit expression of homologous genes in different plant species.

20 In sense suppression, the introduced sequence, requiring less than absolute identity, need not be full length, relative to either the primary transcription product or to fully processed mRNA. A higher identity in a shorter than full length sequence compensates for a longer sequence with less identity. Furthermore, the introduced sequence need not have the same intron or exon
25 pattern, and identity of non-coding segments may be equally effective. Sequences of at least 50 base pairs are preferred, with introduced sequences of greater length being more preferred (see, for example, those methods described by Jorgensen et al., *supra*).

Antisense Suppression

In antisense technology, a nucleic acid segment from the desired plant gene is cloned and operably linked to an expression control region such that the antisense strand of RNA is synthesized. The construct is then transformed into plants and the antisense strand of RNA is produced. In plant cells, it has been shown that antisense RNA inhibits gene expression.

The nucleic acid segment to be introduced in antisense suppression is generally substantially identical to at least a portion of the endogenous gene or genes to be repressed, but need not be identical. The vectors of the present invention therefore can be designed such that the inhibitory effect applies to other proteins within a family of genes exhibiting homology or substantial homology to the target gene. Segments from a gene from one plant can be used, for example, directly to inhibit expression of homologous genes in different plant species.

The introduced sequence also need not be full length relative to either the primary transcription product or to fully processed mRNA. Generally, higher homology can be used to compensate for the use of a shorter sequence. Moreover, the introduced sequence need not have the same intron or exon pattern, and homology of non-coding segments will be equally effective. In general, such an antisense sequence will usually be at least 15 base pairs, preferably about 15-200 base pairs, and more preferably 200-2,000 base pairs in length or greater. The antisense sequence may be complementary to all or a portion of the gene to be suppressed and as appreciated by those skilled in the art, the particular site or sites to which the antisense sequence binds as well as the length of the antisense sequence will vary, depending upon the degree of inhibition desired and the uniqueness of the antisense sequence. A transcriptional construct expressing a plant A-type response regulator (e.g., those described herein) antisense nucleotide sequence includes, in the direction of transcription, a promoter, the sequence coding for the antisense RNA on the sense strand, and a transcriptional termination region. Antisense sequences may be constructed and expressed as described, for example, in van der Krol et al. (*Gene* 72: 45-50, 1988); Rodermel

et al. (*Cell* 55: 673-681, 1988); Mol et al. (*FEBS Lett.* 268: 427-430, 1990); Weigel and Nilsson (*Nature* 377: 495-500, 1995); Cheung et al., (*Cell* 82: 383-393, 1995); and Shewmaker et al. (U.S. Patent No. 5,107,065).

In one working example, antisense expression of the gene encoding A-type response regulators of *Arabidopsis* or an A-type response regulator homolog is used to increase plant yield. In one particular approach, a plant expression vector is constructed that contains the cDNA sequence of A-type response regulator homolog in antisense orientation that is constitutively expressed under the control of the rice actin promoter described by Wu et al. (WO 91/09948). This expression vector is then used to transform rice plants according to conventional methods, for example, using the methods described in Hiei et al. (*Plant J.* 6:271-282, 1994).

Dominant Negatives

Transgenic plants expressing a transgene encoding a dominant negative gene product of an A-type response regulator are assayed in artificial environments or in the field to demonstrate that the transgene increases yield, delays senescence, increases photosynthesis, or increases leaf formation on the plant expressing the gene. Dominant negative transgenes are constructed according to methods known in the art. Typically, a dominant negative gene encodes a mutant A-type response regulator which, when overexpressed, disrupts the activity of the wild type polypeptide.

Mutants

Plants having decreased expression of an A-type response regulator are also generated using standard mutagenesis methodologies. Such mutagenesis methods include, without limitation, treatment of seeds with ethyl methylsulfate, ethylmethylsulfonate (EMS), or fast neutron irradiation, as well as T-DNA insertion methodologies. Expression of an A-type response regulator and increased yield phenotypes in mutated and non-mutated lines are evaluated according to standard procedures (for example, those described herein). Mutated

plants having decreased expression of a gene encoding a A-type response regulator exhibit increased yield relative to their non-mutated counterparts.

Use

- 5 The invention described herein is useful for a variety of agricultural and commercial purposes including, but not limited to, improving and enhancing photosynthesis, promoting cell proliferation, shoot meristem formation, promoting leaf development, increasing crop yields, improving crop and ornamental quality, and reducing agricultural production costs. In particular,
- 10 ectopic expression of a B-type response regulator (such as ARR2 or an ARR2 homolog) in a plant cell provides a method for increasing plant productivity, photosynthesis, and delaying senescence. The invention is especially useful for crop plants such as tomato, potato, cotton, pepper, maize, wheat, rice, and legumes such as soybean.
- 15 All publications and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each independent publication or patent application was specifically and individually indicated to be incorporated by reference.
- 20 What is claimed is:

CLAIMS

1. A method for increasing yield in a plant, said method comprising the steps of:

5 (a) introducing into plant cells a transgene comprising DNA encoding a B-type response regulator operably linked to a promoter functional in plant cells to yield transformed plant cells; and

(b) regenerating a plant from said transformed cells, wherein said B-type response regulator is expressed in the cells of said transgenic plant, thereby increasing yield in said plant.

10

2. The method of claim 1, wherein said B-type response regulator is a crucifer B-type response regulator.

15 3. The method of claim 2, wherein said crucifer B-type response regulator is selected from the group consisting of ARR1, ARR2, and ARR10.

20 4. The method of claim 1, wherein said DNA encoding said B-type response regulator is constitutively expressed, inducibly expressed, expressed in a cell-specific, tissue-specific, or organ-specific manner, or expressed under cycling conditions.

5. A method for increasing yield in a plant, said method comprising the steps of:

25 (a) introducing into plant cells a transgene operably linked to a promoter functional in plant cells to yield transformed plant cells; and

(b) regenerating a plant from said transformed cells, wherein expression of said transgene reduces expression of an A-type response regulator in the cells of said plant, thereby increasing yield in said plant.

6. The method of claim 5, wherein said A-type response regulator is a crucifer A-type response regulator.

7. The method of claim 6, wherein said crucifer A-type response
5 regulator is selected from the group consisting of ARR4, ARR5, ARR6, and
ARR7.

8. The method of claim 5, wherein said transgene expresses antisense A-
type response regulator RNA.
10

9. The method of claim 5, wherein said transgene expresses a dominant
negative A-type response regulator.

10. The method of claim 5, wherein said transgene co-suppresses
15 expression of A-type response regulator.

11. The method of claim 5, wherein said DNA encoding said transgene is
constitutively expressed, inducibly expressed, expressed in a cell-specific, tissue-
specific, or organ-specific manner, or expressed under cycling conditions.
20

12. A method for increasing shoot formation in a plant, said method
comprising the steps of:

(a) introducing into plant cells a transgene comprising DNA encoding a B-
type response regulator operably linked to a promoter functional in plant cells to
25 yield transformed plant cells; and

(b) regenerating a plant from said transformed cells, wherein said B-type
response regulator is expressed in the cells of said plant, thereby increasing shoot
formation in said plant.

13. The method of claim 12, wherein said B-type response regulator is a crucifer B-type response regulator.

14. The method of claim 13, wherein said crucifer B-type response
5 regulator is selected from the group consisting of ARR1, ARR2, and ARR10.

15. The method of claim 12, wherein said DNA encoding said B-type response regulator is constitutively expressed, inducibly expressed, expressed in a cell-specific, tissue-specific, or organ-specific manner, or expressed under
10 cycling conditions.

16. A method for increasing shoot formation in a plant, said method comprising the steps of:

(a) introducing into plant cells a transgene operably linked to a promoter
15 functional in plant cells to yield transformed plant cells; and

(b) regenerating a plant from said transformed cells, wherein expression of said transgene reduces expression of an A-type response regulator in the cells of said plant, thereby increasing shoot formation in said plant.

17. The method of claim 16, wherein said A-type response regulator is a crucifer A-type response regulator.

18. The method of claim 17, wherein said crucifer A-type response regulator is selected from the group consisting of ARR4, ARR5, ARR6, and
25 ARR7.

19. The method of claim 16, wherein said transgene expresses antisense A-type response regulator RNA.

20. The method of claim 16, wherein said transgene expresses a dominant negative A-type response regulator.

21. The method of claim 16, wherein said transgene co-suppresses
5 expression of A-type response regulator.

22. The method of claim 16, wherein said transgene is constitutively expressed, inducibly expressed, expressed in a cell-specific, tissue-specific, or organ-specific manner, or expressed under cycling conditions.

10

23. A method for delaying senescence in a plant, said method comprising the steps of:

(a) introducing into plant cells a transgene comprising DNA encoding a B-type response regulator operably linked to a promoter functional in plant cells to
15 yield transformed plant cells; and

(b) regenerating a plant from said transformed cells, wherein said B-type response regulator is expressed in the cells of said plant, thereby delaying senescence in said plant.

20 24. The method of claim 23, wherein said B-type response regulator is a crucifer B-type response regulator.

25 25. The method of claim 24, wherein said crucifer B-type response regulator is selected from the group consisting of ARR1, ARR2, and ARR10.

26. The method of claim 23, wherein said DNA encoding said B-type response regulator is constitutively expressed, inducibly expressed, expressed in a cell-specific, tissue-specific, or organ-specific manner, or expressed under cycling conditions.

30

27. A method for delaying senescence in a plant, said method comprising the steps of:

(a) introducing into plant cells a transgene operably linked to a promoter functional in plant cells to yield transformed plant cells; and

5 (b) regenerating a plant from said transformed cells, wherein expression of said transgene reduces expression of an A-type response regulator in the cells of said plant, thereby delaying senescence in said plant.

28. The method of claim 27, wherein said A-type response regulator is a
10 crucifer A-type response regulator.

29. The method of claim 28, wherein said crucifer A-type response regulator is selected from the group consisting of ARR4, ARR5, ARR6, and
15 ARR7.

30. The method of claim 27, wherein said transgene expresses antisense A-type response regulator RNA.

31. The method of claim 27, wherein said transgene expresses a dominant
20 negative A-type response regulator.

32. The method of claim 27, wherein said transgene co-suppresses expression of A-type response regulator.

25 33. The method of claim 27, wherein said transgene is constitutively expressed, inducibly expressed, expressed in a cell-specific, tissue-specific, or organ-specific manner, or expressed under cycling conditions.

34. A method for increasing yield in a plant, said method comprising the steps of:

(a) introducing into plant cells a transgene comprising DNA encoding a histidine kinase operably linked to a promoter functional in plant cells to yield
5 transformed plant cells; and

(b) regenerating a plant from said transformed cells, wherein histidine kinase is expressed in the cells of said plant, thereby increasing yield in said plant.

10 35. The method of claim 34, wherein said histidine kinase is a crucifer histidine kinase.

36. The method of claim 35, wherein said crucifer histidine kinase is CKI1 or CRE1.

15

37. The method of claim 34, wherein said DNA encoding said histidine kinase is constitutively expressed, inducibly expressed, expressed in a cell-specific, tissue-specific, or organ-specific manner, or expressed under cycling conditions.

20

38. A method for increasing shoot formation in a plant, said method comprising the steps of:

(a) introducing into plant cells a transgene comprising DNA encoding a histidine kinase operably linked to a promoter functional in plant cells to yield
25 transformed plant cells; and

(b) regenerating a plant from said transformed cells, wherein histidine kinase is expressed in the cells of said plant, thereby increasing shoot formation in said plant.

39. The method of claim 38, wherein said histidine kinase is a crucifer histidine kinase.

40. The method of claim 39, wherein said crucifer histidine kinase is
5 CKI1 or CRE1.

41. The method of claim 38, wherein said DNA encoding said histidine kinase is constitutively expressed, inducibly expressed, expressed in a cell-specific, tissue-specific, or organ-specific manner, or expressed under cycling
10 conditions.

42. A method for delaying senescence in a plant, said method comprising the steps of:

(a) introducing into plant cells a transgene comprising DNA encoding a
15 histidine kinase operably linked to a promoter functional in plant cells to yield transformed plant cells; and

(b) regenerating a plant from said transformed cells, wherein histidine kinase is expressed in the cells of said plant, thereby delaying senescence in said plant.

20

43. The method of claim 42, wherein said histidine kinase is a crucifer histidine kinase.

44. The method of claim 43, wherein said crucifer histidine kinase is
25 CKI1 or CRE1.

45. The method of claim 42, wherein said DNA encoding said histidine kinase is constitutively expressed, inducibly expressed, expressed in a cell-specific, tissue-specific, or organ-specific manner, or expressed under cycling
30 conditions.

46. A plant or plant component comprising at least one transgene encoding (i) an A-type response regulator polypeptide, (ii) an antisense A-type response regulator RNA, (iii) a dominant negative A-type response regulator polypeptide, (iv) a B-type response regulator polypeptide, (v) an antisense B-type response regulator RNA, (vi) a dominant negative B-type response regulator, (vii) an antisense HK RNA, (viii) a dominant negative HK polypeptide, or any combination of (i) –(viii).

47. The plant or plant component of claim 48, further comprising a transgene encoding a histidine kinase polypeptide.

48. A plant or plant component comprising:

(a) a first transgene encoding (i) an A-type response regulator polypeptide, (ii) an antisense A-type response regulator RNA, or (iii) a dominant negative A-type response regulator polypeptide or combination thereof;

(b) a second transgene encoding (iv) a B-type response regulator polypeptide, (v) an antisense B-type response regulator RNA, (vi) a dominant negative B-type response regulator or combination thereof, and

(c) a third transgene encoding (vii) a HK polypeptide; (viii) an antisense HK RNA, (ix) a dominant negative HK polypeptide or combination thereof.

49. The plant or plant component of claims 46, 47, or 48, wherein said plant or plant component is selected from the group consisting of wheat, rice, maize, barley, potato, soybean, tomato, oats, cotton, and sunflower.

FIG. 1A




FIG. 1B




FIG. 1C




FIG. 1D




FIG. 1E





FIG. 1F



5/9

FIG. 2A




FIG. 2B




FIG. 3A





FIG. 3B



7/9

FIG. 3C




FIG. 4A




FIG. 4B




FIG. 4C





FIG. 5



SEQUENCE LISTING

<110> The General Hospital Corporation

<120> Cytokinin Response Regulators and Uses
Thereof

<130> 00786/402W02

<150> US 60/296,554
<151> 2001-06-06

<160> 30

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 2401

<212> DNA

<213> Arabidopsis thaliana

<400> 1

tagatcaagt tttaacaatt tttaacgta cttaaaagat tgataaaat gtaactgttg 60

gataaaat agcttttgca acccttttat cactaaaact agcttttgca acccttttat 120

ttgttgacca ttgttgatg tatgttttta ttatcttaact ctatttaatt atctcatgtag 180

cttaacatat caataataac taacatgcga taatctcga agaagaaaaa ttgttataaa 240

agaactatat gcaactaaca aaaaaccaa aaaaaccaa aaaaaccaa ttgttggttaca 300

taaaaaatca ttgttatgaa gaagaaagaa gaatcattac acagctttca ctcaacaagt 360

tacaatgcat ataaggaac tcccaaaaaa tatcattcaa gatctttca 420

gaacaaaaaa aaaaaggtt aaaaatatat ctctttgccc atagttagac tcaaaaaaac 480

gacatcgttt accccaacga tgcacaacta cgcacttaca gaagaaacca caagtacytt 540

gaaggtaact tagtcaattta gcatcgaacg aaatgtcttg agcggaggaag aggtatattcc 600

gttatccctt tatatatat ctctctcacc ttctctcctt ttctcttgat ttttataa 660

atcaaaaaaa atagaaaaaa aaaaataaaa ataaaaataa aaaaatctca cgtttcttct 720

ctctctcctc ctctctcctc cgaagccaac aatcttgaaat aggttttttg agaatattca 780

tcatttgat tcgaattctt caccacatgt gtaattcaac tctgtcaggt tcatctgcgg 840

aatcatgat acagattcga tcaaccaacg ttctgtcact tcagctccga gagattccgg 900

aaagaagaag aggttaacgt tatccctctt gatataatctc atcccttctc ctgaatatga 960

ttcaaaattt tgatttttaa tggttttgtt attgcattac gtttttgagga acaataaagtc 1020

ggctaagatg aagcagaaca agcttggctt caatggcttt ctcaaggtta 1080

aaacttcatc ttctctcctt ttaattgttc aatctcgtct gatctglttg gttagatttg 1140

gtttgatctt tatgtatttt ggttttttga tttrtgatat agttgaggtg agcaataaag 1200

aaagttaaga ggaagagaggt gttaatcgtat gttcaaaagcc tcatctagag agttcagata 1260

aggttcgtag aagaagaggtt aacaatgttg ggaatattct tcttcatcat gagagtatta 1320

tggaagtcaacc ttcaatatagc tctgtttgtg gtaatatatc gagcaactaac ttcagtggga 1380

gaagttagcag gagttagat agcagcagtg gctttgtctc tggtaatatc acagaagagg 1440

aaaatgttaga cgtatgtgat gatgggttg gatgtgatg ggaagcgttt gctgatgtcg 1500

tagcggctga ggaagagat gagaaaaaga gtcgtcctct tgaagtcgtg aaagagcaag 1560

tgaagtcgtg acaatcagct tctaattgtg ttgtatctgc gatcaggtgat gcatcagaag 1620

ttgttggttg tgaagatcca aagcaggaat gttcagaggt gttcagaggt gttcagaggt 1680

gtaatatagag ttgtaggcta gatgtgacc ttgcaccaaa ggggttaact aatttgcga 1740

agcagcttag tttrccggag ttagaacaag gtttttagct tgttgcgatc cgttctcat 1800

gtcccatag ctacgaagac ttggaactga cggatcgaat ttccctcccc tgtcctgttg 1860

gattcggct cgtctcgttc tgcacaaga ccatctgcga ttggagatggg cgtgtccag 1920

gctgcaggaa accctatgaa cggatatgg tcaaggtcga gactagttat caaggtgttg 1980

gtctaaacaat tcggttggt cgttcgtcta gcatgttttg caagttttaa aagggagggt 2040

goggttttct caacatgtt gttcttttga actcagaaac tttaagctctg ttttctatgt 2100

```

ttgaaaggag tgactcacgg tgcagttgat tacctcatca aaccgggtacg tattgaggct 420
ttgaagaata tatggcaaca tgtggtgagg aagaagcgta acgagtggaa tgtttctgaa 480
cattctggag gaagtattga agatactggc ggtgacaggg acaggcagca gcagcatagg 540
gaggatgctg ataacaactc gtcttcagtt aatgaaggga acgggaggag ctcgaggaag 600
cggaaggaa aggaagtaga tgatcaaggg gatgataagg aagactcatc gagtttaaag 660
aaaccacgag tggtttggtc tgttgaattg catcagcagt ttgttgctgc tgtgaatcag 720
ctaggcggtg acagtgagtt aaaaacttgc ttgcttatgc atttgtgtgt gtcgattggt 780
aacattgtgg aattccagaa gtatcggata tatctgagac ggcttggagg agtatcgcaa 840
caccaaggaa atatgaacca ttcgtttatg actggtcaag atcagagttt tggacctctt 900
tcttcgttga atggatttga tcttcaatct ttagctgtta ctggtcagct ccctcctcag 960
agccttgcac agcttcaagc agctggtcct ggccggccta cactcgctaa accagggatg 1020
tcggtttctc cccttgtaga tcagagaagc atcttcaact ttgaaaaccc aaaaataaga 1080
tttgagagac gacatggtca gacgatgaac aatggaaatt tgcttcatgg tgtcccaacg 1140
ggtagtcaca tgcgtctgag tcctggacag aatgttcaga gcagcggaaat gatgttgcca 1200
gtagcagacc agctacctcg aggaggacca tcgatgctac catccctcgg gcaacagccg 1260
atattgtcaa gcagcgtttc aagaagaagc gatctcactg gtgctgctgg ggttagaaac 1320
agtatccccg agaccaacag cagagtgtta ccaactactc actcggctct caataacttc 1380
cccgcggatc tacctcgag cagcttcccg ttggcaagtg ccccgaggat ttcagttcca 1440
gtatcagttt cttaccaaga agaggtcaac agctcggatg caaaaaggag ttcatcagct 1500
gctactgctg gatttggtta cccaagctac gacatattta acgattttcc gcagcaccaa 1560
cagcacaaca agaacatcag caataaacta aacgattggg atctgcggaa tatgggattg 1620
gtcttcagtt ccaatcagga cgcagcaact gcaaccgcaa ccgcagcatt tccacttcg 1680
gaagcatact ctctgctctc tacgcagaga aaaagacggg aaacggacgc aacagttgtg 1740
ggtgagcatg ggcagaacct gcagtcaccg agccggaatc tgtatcatct gaaccacgtt 1800
tttatggagc gtggttcagt cagagtgaag tcagaaagag tggcggagac agtgacttgt 1860
cctccagcaa atacattgtt tcacgagcag tataatcaag aagatctgat gagcgcattt 1920
ctcaaacagg tttga

```

<210> 4

<211> 599

<212> DNA

<213> Artificial Sequence

<220>

<223> derived from *Arabidopsis thaliana*

<400> 4

```

gctttgaaga atatatggca acatgtggtg cggaagaagc gtaacgagtg gaatgtttct 60
gaacattctg gaggaagtat tgaagatact ggccggtgaca gggacaggca gcagcagcat 120
agggaggatg ctgataacaa ctctgtcttca gttaaatgaag ggaacgggag gagctcgagg 180
aagcgggaagg aagaggaagt agatgatcaa ggggatgata aggaagactc atcgagttta 240
aagaaaccac gctgtggttg gtctgttgaa ttgcatcagc agtttggtgc tgcgttgaat 300
cagctaggcg ttgacagtga gttaaaaact tgcttgctta tgcattttgt tgtgtcgatg 360
tttgttttct tgtctaactc ttaccgtatt ggggcattca atgcagaagc tgttcctaag 420
aagatccttag agatgatgaa tgtaccggg ctaacgcgag aaaacgtagc cagtcacctc 480
caggatatata tgtgatcgaa tagtattaca ttttttcatt gttgatgtgt tctaacagt 540
gtaacattgt ggaattccag aagtatcgga tatatctgag acggcttgga ggagtatcg 599

```

<210> 5

<211> 780

<212> DNA

<213> *Arabidopsis thaliana*

<400> 5

```

atggccagag acggtggtgt ttcttgttta cgaaggctcg agatgatgag cgtcgggtgt 60
atcggaggaa ttgaatctgc gccgttggtt ttagatgaag ttcattgtct agccgttgat 120
gacagtctcg ttgatcgat tgtcatcgag agattgcttc gtattacttc ctgcaaagt 180
acggcggtag atagtggatg gcgtgctctg gaatttctag ggtagataa tgagaaagct 240

```

<212> DNA
<213> Arabidopsis thaliana

<400> 8

caatctgctc tctttttat tctgagttg acaatgacgg ttggtgaggt catgagagtg 60
gagatctccg ccggtggaga ttgactgtt actactccgg agttgcatgt tcttgccgtc 120
gtatgagta ttggtgacg taaagtcatc gtaggtgtgc ttagaatctc tcttgttaa 180
gtacgacctg atctaaagga ttgaaaggtg aactagggct ttggccttag ttgaggtcaa 240
ggggcttcta atctaaagga ttgaaaggtg aattgtag ttgaggtatc ttgaggtcaa 300
ggactttcag gatattgact ccttaaaaaa attaaaggaat cttcagcatc cagaggaagt 360
ccagtatgta ttatgtcatc tgaagaacac taccctcgta tacaagaaatg tctcaagaag 420
ggagcagagga aattctgtt gaaacccgtg aagcttagcag atgtaaagcg aatcaacaa 480
ccttaataga ggaatgaagc tgaagaaatgc aaaaatctaa gccattctaa caagagaagt 540
ctcaagaaag acagtgtac atcatcatca agtcatgtag ttctctact ttgaattata 600
tcatgttcaa aacgaatga atcagaatc gaaaacctt ttctctact ttgaattata 660
ttgtatagag ccaagacct agcaaaaacca ttctgttaat attcctcag aagacaatg 720
tttgaagag tcatcacact

<210> 9
<211> 1659
<212> DNA
<213> Arabidopsis thaliana

<400> 9

atgactatgg agcaagaaat tgaagtcttg gaccagtctc cggttgggat gtagttctt 60
gctgtltgacg atgaccagac ttgtctccgt attctccaga ctttgcttca gctgtgccaa 120
taccagctta caaacagaa agttgtttga agttgtttga gctgtgccaa ctttgcttca 180
gagagaaagc ctcgtgtctc ttggaagcat ttggaagcat ttggaagcat ttggaagcat 240
gaaagaaagga agagaaagga agagaaagga agagaaagga agagaaagga agagaaagga 300
aatagaagca atgtgagtaa ttgttcagga aactgtgata aagcaaacag aaaaagtaaa 480
cgaattgagg agcttaagaa catattggcaa catattggcaa catattggcaa catattggcaa 540
gagagaaagt agctgttcca agctgttcca aaaaagatc ttgagaaat tccggttg 720
aaactcaata gagagaaagt ttggaagcat ttggaagcat ttggaagcat ttggaagcat 780
gtgtcttgat accgactca acaagctaac agggcggcta ttgactcaaa ttttatgaa 840
atgaattctc agaaaggaat ttgtgtgtc taccacacac accgctggaat accgtgtgga 900
tccgttcagt tccatgtgtg aacacaatg atgaggtcat actcttcaa taggaatct 960
aaccatcatc acaaggcttt tccaagctt acttcaaac aaaaactctc aatggttagc 1020
ccaagtaatc tgttactct cgaaggtaac ccgaggtcat catcttacc ctcaaacccg 1140
ggtttttctc ctcatctcga gatcagcaag cgtctagaac attgttcaaa cgtgtcatg 1200
tcaacaaaca tccacagag ttgattctat tcaaaacctg acaaccttga atgtgaatgcg 1260
ttctgagact cagctagccc gctagttaac ccaaaccttg atacaatcc ggcatctctc 1320
tgcagaaaca cgggttttgg atccaacaat gctgcacaaa cagaactctc ttatccata 1380
cagatgatatc agaacctgc aaaaacatca ggtccaagtga agtccaagag agcttcagag 1440
agttagcaatc caaacgaag tttaactag gtaactcatg ggaacagagc ttggtttag 1500
gcttctgagt ctgttccctt agatgtatat ttgattggat gtaattggat ggaacagagc 1560
caatctgact tctcggaaag ttgattggat ttgattggat ttgattggat ttgattggat 1620
tgcagagaaac tccatttcc ctttctctg tcagtctga

<210> 10
<211> 1935
<212> DNA
<213> Arabidopsis thaliana

<210> 12
 <211> 942
 <212> DNA
 <213> Arabidopsis thaliana

<400> 12
 atcctccaac attcgttgat caatggctga agttatgcta ccgaggaaga tggagattct 60
 caaccattct tcaaagtttg gatcaccgga tcctcttcat gttcttgccg tcgacgacag 120
 tcacgttgat cgtaaattca tcgagcggtt gctcagagta tcttcttgca aagttactgt 180
 tgttgatagt gcaacaagag ctctccaata ccttggtttg gatgttgagg aaaaatcagt 240
 cggttttgag gatttgaagg ttaatttaat tatgactaat tactcaatgc ccggtatgac 300
 tggatatgaa ctcttgaaga agatcaaaga atcctcagct tttagagaag taccggtggt 360
 aattatgtcc tccgagaaca ttttgccctcg tattgataga tgtcttgaag aaggggctga 420
 agatttctta ctgaagccgg taaaactctc ggatgtaaaa agattaagag attctctaatt 480
 gaagggtgaa gatttgtctt tcacaaagag tattcagaag agagagctag aaacagagaa 540
 tgtctaccct gttcactcgc agctcaaagc cgcaaagatc tgagctctcc gatgcaaatt 600
 ccgtgactgg atcttagaaa aagccttttc atatacagga agcttcaaga ctcaatgcac 660
 ttgagctaca ctgatgattc tcgccagctt aatgacctat acaattttcc aaccctatg 720
 gcccttttctg ctgattgaat tcagtgggtt tattcatata gtagtaattgt aaatgggtata 780
 gagatgagca aaagctatat tatatattat attcattaga ggggtcatag cttttttcct 840
 tacaagagat ttgtaagctg gcttaataga agttgttaag aagcctcact ttatgtgaat 900
 aatgattttt gttagctaatt cataaacaat cagaagtttc ct 942

<210> 13
 <211> 3677
 <212> DNA
 <213> Arabidopsis thaliana

<400> 13
 aattattcac tcaaattcac aaagggttcc gactttgctc cgaggaagaa gataatatga 60
 aaagagcttt ttaggggttta tcattctcct tgactttgca aaacgtgaaa taagcttctt 120
 ccttgaggga ctaaatcaaga cagaaatctg ttctctctaa aacgatcgcc gttctagttc 180
 actcaaatga tgggtgaaagt tacaaagctt gtggcttcac gtccaattgt ggtcttttgc 240
 gtcttgccat tcttggtggt tgttttcgag tgcatttggg tctcaaattg gcgaacaaca 300
 acggagaacc tagtcaaaga ggtcgcttca tttaccgaag atctccggac aagtctagtt 360
 tcggagattg aaaacatcgg aaaatttaca tatgctaaga caaacttate tacgatcggg 420
 ttagcgagag ttatagattc ttatatcacc aacaacgaca ctgggttttac agagattcaa 480
 acacagatcg caccattggt gttttagctt tattcaacga tccttcaagt ctcaaaagtt 540
 tcgtacatca gtatgggacgg tctcatgttt tcttacattg cagaatcaaa cacaagtgtc 600
 gctgtttttg ccaattcctc gtcgaaattca agtcgtggag actacacttg gtacactcaa 660
 accgtggatc agttaactgg tcgtcttaac ggggaactcaa cgaaatctca gtcgttagat 720
 gtaaccata cagattgggt ccaagcagca cagagtaata actacactac agcctttgta 780
 ggaacgagct tgggaggaga agataacgag actctaatac agagcgtggt tagcttgtac 840
 agcaagaaag gtcttggttc tttaggggtt ccggttaaga ctttaaccga agttttgaac 900
 agtttgaate tacacggcga agagctttac atgtggacaa aggacgggac ggtgcttggt 960
 cgtgaagggt cactgaatga ttctttcttc cctgaaaatt gcagttccag tggctacgag 1080
 gaatcgaact cctctgggtc tcaatgcata cctgaaaatt gcagttccag tggctacgag 1080
 gtggagatca aaagattaag ataccaagct ttttgcctg ttattgaagt ttccggcggt 1140
 cctctgagat acacactcat gtttcccaac aaaggaggag caacacgcat caagcaccia 1200
 gcggaaaagg caaaatatca acttattgtg gttatgatat ttcttggtt cggttggcct 1260
 gtatggtttg tgtgggtttat gatgcaagca acaaggagag agatgcatat gcgtgcaacg 1320
 ctgataaacc aaatggaagc gacacaacaa gctgagagaa agagcatgaa caagagtcaa 1380
 cgatttgcaa atgctagcca cgatattaga ggtgcccttg cagggatgaa aggtctgatt 1440
 gatatatgtc gtgatggagt taaacctggc tccgacgtag acaccactct caaccaagt 1500
 aatgtttgag ccaaggattt ggttgcctc ctcaactctg ttttggacat gagcaaaate 1560
 gaaagcggga agatgcagtt agtgaagaa gatttcaact tgtcgaaact tcttgaagac 1620
 gtcatcgatt ttaccatcc cgttgcgatg aagaaagggg ttgatgtagt tttggatccg 1680
 cagcatggct cggttttcaa attctcgaat gtacgagggg atagtggcag actgaagcag 1740

agtttgatc tacacggcga agagcttac agtggaacaa aggaaggga ggtgcttggt 960
 cgtgaagggt cactgaatga ttctttcttc atctccaatg gctcgaattg cttcggtaga 1020
 gaatcgaact cctccttggtc tcaatgcatc cctgaataat gcaggtccag 1080
 gtbgaagatca aagatgaag ttctggtcctg ttatbtgaagt ttccgggctgt 1140
 cctctgaagt acaactcat gttcccaac aaaggagag caacacgcat 1200
 gccgaataaagg caaatatca acttatgtg gttatgatbt ttcttggtc 1260
 gtabgttttg ttgtggttat gttgcaagca acatgaagag agatgcatga 1320
 ctgataaac caatggaag gacacacaa gttgaagaa gttggtccctg 1380
 gcatbtgcaa atgtcatgaa agtatatga ttccgacttg caggatbga 1440
 gatbtatgtc gttgatbga ttcaaaccttg ttccgacttg acacacactc 1500
 aatgttttgcc caaggatct gttgtgcttg ttcttggaat ttcttggaat 1560
 gaaagcggga agatgcagtt gatttcaact ttgtcgaact ttcttggaat 1620
 gtcactgat tttaacctc cgttgcgatg aagaaagggt ttgatgtagt 1680
 cactgatbgtc atctcgaat ttctcgaat gttacgggggt atagtgggca 1740
 atctgaaca atctgttag caatgctgtc aagttcaccg ttcaagggtca 1800
 agagcttggt ctcaggagcc aggttccaat agctcttggt tccctgcatc 1860
 ggttggtacca agttgtacca ttgcaagaa ttgcaagaa ttgcaagaa 1920
 gagaacagaaa tatcgatcc catgaagaa ttgcaagaa ttgcaagaa 1980
 gttgatgatba ctgttaagg gataacctg agatgtcgta ttcttggtc 2040
 gttcaagtaa gagaacagca ttcaaggac caaggaaactg gtttagggt 2100
 caatgtcttg taagatatg gggagggggg ataaagaaatca ttgaagaa 2160
 gaaagcaact gtttccaat caatgtttta ttgaacaaact ttgaagttgt 2220
 gacatgaaag ttgaacagga gatcggaagca ggaaggcgtt atgtatccac 2280
 gggcttgacta taaacactc acttggaagt agcatgaaata taagtccatga 2340
 ttcaacaact gttcagctc aagttccaaag caagaagttt ctatgatgt 2400
 aaaaatgaag aacgcagaa agttacagag aatatcaata aatatccttg 2460
 acaagtgttg aaaaatggga gcatrttgat tatrgtcttg agagaacttt 2520
 cctcaggagt ccatgggaag agcaggttcat gttcgggtc aagggtgcat 2580
 ccttcatat gcatggaggg tatgatcca agaaagccaac ttccataag 2640
 agtttctctg cagttgtctt ttgtgtgat ttgtgagctg 2700
 ttgcatatctg tcaaacagtt ttgtagaggg ttgttgatccctg 2760
 ttgcttaacg aatcgagcac ttgttgatg gagaagaggg aatatgttg 2820
 ttgcaacggat cgttcttat ggaagtcttg aagatgttg ctgaatttg 2880
 ctaaaagaa cactactga gttgcaaaag gaatcalttg ttgaacatct 2940
 taacaataa gttcgaatg gttcgaatg ttgaacatg ttgaacatg 3000
 tttagtcaagtt caagttcgaatg ttgaacatg ttgaacatg 3060
 tttagtcaagtt caagttcgaatg ttgaacatg ttgaacatg 3120
 gacgatbgaat tttaagag ttctggtctg atgtatcaact ttatcaact 3180
 aaagtgtcaa caggaaagtt ggaagtctcaac ttgaactcaac 3240
 gggaaagaa ctttgagatt agttcaactg ttgaactcaac ttgaactcaac 3300
 gtagatbgaat ttccgttgca ttcaactatc ttgaactcaac ttgaactcaac 3360
 tatgaagcaat ttgaagagtt ttgaagagtt ttgaagagtt ttgaagagtt 3420
 atagctgat atagctgat ttctggtctga ttctggtctga ttctggtctga 3480
 atggagcctc ttctgaataa ttctgaataa ttctgaataa ttctgaataa 3540
 agcaaacgtc actagatctg ttatgaatga ttatgaatga ttatgaatga 3600
 ttgttttagac aatactttt tagtatgacg atgtcatagc agtttcttc 3660
 taaagttcat ttcatgat 3677

<210> 15
 <211> 3677
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> derived from Arabidopsis thaliana

<400> 15
 aattatcac tcaaatcac aaagggttc gacttgctc cgaggagaa gatataatga 60

```

agcaaacgtc actagtattg tataagtaga cagcacgcag ctttgtgcta ctaccattag 3600
tgttttagac aatacttttt tagtatgacg atgtcatagc agttttcttc taagttttatt 3660
taaagtcocat ttcagat 3677

```

<210> 16

<211> 3531

<212> DNA

<213> Arabidopsis thaliana

<400> 16

```

atgtctataa ctttgtgagct cttgaatctt acttcaaaga aagctaagaa gtcgtcgagc 60
agtgacaaga aatggctaaa gaagcctctc ttcttcctga ttttgtgtgg ctcttttgga 120
attgttttgg ttatgttctt acggttaggt agaagtcaga aggaggagac agattcttgt 180
aatggagaag agaaagtgtt gtatagacat caaaatgtca caagaagtga gattcatgat 240
ttgggtctctt tgttctctga ttcagatcag gtaacatcct ttgaatgtca taaggaatca 300
agccctggaa tgtggacaaa ctatggtatt acatgttccc tgagtgtgag ttctgataaa 360
caagagacta gagggcttcc ctggaatctt ggcttaggac attctatctc atcaacatct 420
tgtatgtgtg gtaatcttga accgatttta cagcaacctg aaaaccttga ggaagaaaac 480
catgaagaag ggctggagca ggggttgtca tcgtatttaa gaaatgcag gtggtgtcta 540
atccttgggt tgttagtgtg ccataagatt tatgtatctc attctaaagc acgaggtgag 600
aggaaagaga aagtacatct gcaagaggct tttagctcaa agaagcagca acaacgtgct 660
cagacttctt cttagagggc tggaaagtgg aggaagaata tccttctcct tggattttta 720
ggaggagttt ccttctctgt ttggtggttt tgggacacta atgaggagat cataatgaaa 780
aggagggaga ctttggcaaa catgtgtgac gaacgagcac gtgttttaca agatcagttc 840
aatgttagct tgaacctatg tcatgccttg tctattcttg tatctacatt tcatcatggt 900
aaaatcccat ctgccattga tcagagaaca tttgaagaat atactgagag aacaaaacttt 960
gagagggccac ttactagtgg tgtagcgtat gctttgaaag tcccacactc agaaagagag 1020
aaatttgaaa aggagcatgg atgggcaata aagaaaatgg aaactgagga ccagacagtt 1080
gtacaagatt gtgttctga aaactttgat cccgcaccga ttcaagacga atacgcgcca 1140
gttatatttg ctcaagaaac tgtttcccat attgtatcgg tcgacatgat gtctggagaa 1200
gaagaccgtg aaaacatctt acgggcaagg gcatcaggaa aaggagtgtt aacatctcca 1260
tttaagcttc ttaagtcaaa tcatcttggt gttgtgttga cctttgctgt ctatgacacg 1320
agcctaccgc ctgatgctac agaagaacag cgtgttgaag caactattgg gtaccttggt 1380
gcacatcatg atatgccatc gctgggtggag aaactcttc accaacttgc cagcaaacag 1440
acaattgctg tggatgttta cgacacaact aacacttcag gtctaataaa aatgtatggc 1500
tcagaaattg gggatataag tgagcagcat ataagtagcc ttgattttgg tgatccatca 1560
aggaaccatg agatgcattg caggtttaag cataaacttc ccattccctg gacagcgata 1620
acaccgtcga tcttagttct gggtattact tttcttgggt gttatatattt atatgaagcc 1680
atcaaccgaa ttgcgacagt tgaagaggat tgtcagaaga tgagggaaact caaagctcgt 1740
gctgagggcg ctgacattgc aaagtcacag ttcttagcaa ctgtttctca tgagatacgg 1800
actccgatga atggagtttt aggaatgctg aaaatgctga tggacaccga tcttgatgag 1860
aagcagatgg actatgcgca aactgctcat ggcagtggga aggatcttac atcactaata 1920
aatgaggttc ttgatcaggc aaagattgaa tccggaaggc tcgagcttga aaatgtgcct 1980
tttgatatgc gttttattct tgataatgtt tcatctctcc tctctggcaa ggcaaatgaa 2040
aaaggaattg agttggccgt ttatgtttct agtcaagttc ctgatgttgt agtcggtgat 2100
ccgagtcggt tccggcagat cattacaaac ctggttgga actcaatcaa attcacacag 2160
gaaaggggac acataatttat ctcatgacac cttgcagatg aggtaaagga gcctcttact 2220
attgaagacg cagtgtctaaa acagcgacta gctttaggat gcagcgagtc cggtgagaca 2280
gttagcgggt ttctgcggtt aaatgcatgg ggaagctgga agaatttcaa gacatgttac 2340
agtactgaga gtcagaattc tgatcaaate aaattgctag ttacagtgga ggacactgga 2400
gttggcatac ctgtggatgc acaaggccga atcttcacac cttttatgca agccgacagt 2460
tcacatcgc ggacttatgg tggaaactggc ataggtttga gtataagcaa acgtttgggt 2520
gaactcatgc aaggagagat ggggtttgtg agtgagcccg ggataggcag tactttttca 2580
ttactggag ttttcgggaa agcagaaaca aatacgtcga ttactaagct ggaacgattc 2640
gatctagcta ttcaggagtt tacaggattg agagcattag ttattgataa cagaaacatt 2700
agagcagagg tcaccaggta cgaacttcgg agactgggaa tatctgcaga cattgtttca 2760
agtctgagaa tggcatgcac ttgtgtatc agcaaattag aaaatttggc tatgattcta 2820
atagacaaag acgcctggaa caaggaagaa ttttcagtac ttgacgagtt gtttaccgaa 2880

```

gaagttcttg atcgcgcgaagc gattggaagct ggaagagctgg agttggaaatc agtaaccattt 300
gatataccgtt caatatltgga tgaattccctt tctctcattct ctgaaggagtc aaggaacaataa 360
agcatltgagc tccgcgttttt cgtttcagac aagtaaccag agatagttcaa aggaagtattca 420
gggaagtattt gacagataat catataaccctt gttggaaatt cggtttaatt cacagaagaata 480
ggacatatct ttgtttaagt ccatcttgcg gaaacaatcaa aagatgaaatc tgaaccggaata 540
aatgcatlga atgttggaat caatgttagg ctatatgttt caatcgaaga caagggtattt 600
ggaatccctt tagtttgcgca aggcgttggt ttatgtccgt ttatgtccagc agatcagctcg 660
acctggaagg acctggaagg tactgggtat ggttttgatg taagccaagtg taagccaagt 720
actatgcgtg gtcaagataa ttctataagc cggcctcata ttggaaagcac gttcctgttcc 780
acggcgtgtt tagaagaatg cgaataatgc agtgcgattt accatatgaa gaaaccattat 840
gtggaaacact tgccttctac tttaaaagg attgaaagct tagttgttga ttgctaagcct 900
gttagagctg ctgtgactag ataccatatg aaaaagactcg gaatcaatgt ttgatgtcgtg 960
acaagttcga aacccgctgt tgttgagct gtcgcgtttg aaagaaacgg ttctcctctc 1020
ccaacaataa ccgaactlga tatgaactta gtagagaagc attcatlgtgatt tccaactlga 1080
gataatlgact tagctctatt cgcacaataa atccaataa acgggaacga acgggaacga 1140
tctccgaac tagctctatt cgcacaataa atccaataa acgggaacga acgggaacga 1200
tccgaagat ttgcagatag ggtataatg aaaccgttaa gagcaagcat gattggggcg 1260
tgctcgaac aagttctcga gctggaataa aacaagacaac aacatccaga aggatccata 1320
cccgcaactc tcaagagctt gctcaaggg aagaagattc gagcaagaat ggtttgtgca 1440
gagatgtgttc aagtttgctt gggtttgctt cagatccac acactttcga tgccttgcttc 1500
atggatatcc aatggacca gaattgaga aggaactaa agttgataca cgcgaactac 1560
aatctcgaaat ggactttaac gatctcagc atgactgcgg agttgataca cgcgaactac 1620
gaggaatgtc tgaagaagtgg gatggatgggt tactgtcca aacctttga agaaagaaat 1680
ctctataaat cggttgcca atcatlcaaa ctaatccca ctaatccca cgggaagagt 1740
ggaccgatcc tgcgtatga tagcatttt gtttctgta catcagtag gttccaatc 1800
tag

<210> 19

<211> 1803

<212> DNA

<213> Artificial Sequence

<220>

<223> derived from Arabidopsis thaliana

<400> 19

atgcacatag taaaaagtcga agatgatltc catgaaatgc aagagcttaa agttcggagca 60
gaaagctcgt atgtcgcata atcgcagttt ctgtcatacg ttgtctcaag gatcagga 120
ccaatgaatg gcatctcgg aatgcttgct atgctcctag atacaagaat aagctcgaca 180
cagagagatc acgctcaaaa cgtccaagta tgtgttaaaa agttgttgc atttgataat 240
gaggttcttg gacgcgcga gatgaaagc gttgaaagc agttgaaat agttccaatt 300
gatataccgtt caatatltgga tgaattccctt tctctcattct ctgaaggagtc aaggaacaataa 360
agcatltgagc tccgcgttttt cgtttcagac aagtaaccag agatagttcaa aggaagtattca 420
gggaagtattt gacagataat catataaccctt gttggaaatt cggtttaatt cacagaagaata 480
ggacatatct ttgtttaagt ccatcttgcg gaaacaatcaa aagatgaaatc tgaaccggaata 540
aatgcatlga atgttggaat caatgttagg ctatatgttt caatcgaaga caagggtattt 600
ggaatccctt tagtttgcgca aggcgttggt ttatgtccgt ttatgtccagc agatcagctcg 660
acctggaagg acctggaagg tactgggtat ggttttgatg taagccaagtg taagccaagt 720
actatgcgtg gtcaagataa ttctataagc cggcctcata ttggaaagcac gttcctgttcc 780
acggcgtgtt tagaagaatg cgaataatgc agtgcgattt accatatgaa gaaaccattat 840
gtggaaacact tgccttctac tttaaaagg attgaaagct tagttgttga ttgctaagcct 900
gttagagctg ctgtgactag ataccatatg aaaaagactcg gaatcaatgt ttgatgtcgtg 960
acaagttcga aacccgctgt tgttgagct gtcgcgtttg aaagaaacgg ttctcctctc 1020
ccaacaataa ccgaactlga tatgaactta gtagagaagc attcatlgtgatt tccaactlga 1080
gataatlgact tagctctatt cgcacaataa atccaataa acgggaacga acgggaacga 1140
tctccgaac tagctctatt cgcacaataa atccaataa acgggaacga acgggaacga 1200
tccgaagat ttgcagatag ggtataatg aaaccgttaa gagcaagcat gattggggcg 1260
tgctcgaac aagttctcga gctggaataa aacaagacaac aacatccaga aggatccata 1320
cccgcaactc tcaagagctt gctcaaggg aagaagattc gagcaagaat ggtttgtgca 1440
gagatgtgttc aagtttgctt gggtttgctt cagatccac acactttcga tgccttgcttc 1500
atggatatcc aatggacca gaattgaga aggaactaa agttgataca cgcgaactac 1560
aatctcgaaat ggactttaac gatctcagc atgactgcgg agttgataca cgcgaactac 1620
gaggaatgtc tgaagaagtgg gatggatgggt tactgtcca aacctttga agaaagaaat 1680
ctctataaat cggttgcca atcatlcaaa ctaatccca ctaatccca cgggaagagt 1740
ggaccgatcc tgcgtatga tagcatttt gtttctgta catcagtag gttccaatc 1800
tag

```

atggatttgg ttcagaagca gaagagtttg caagattaca ccaaactact cttcttagaa 60
gggatttggg acagccagtt cttgcagctg caacaactac aagatgaaag caatccagat 120
tttgtttcac aagttgtcac actcttcttc caagactctg ataggattct caatgatctc 180
tcactttccc tagatcaaca agttgtagac tttaaaaaag ttgatcccca tgttcacaa 240
ctcaaaggta gcagctccag tataggagca cagagagtta agaagcttg tgttgccttc 300
cgcagcttct gcgagcagca aaatgtcgaa gcagtcata gatgtttgca acaagtgaag 360
caagagtatt atcttgtgaa aaacagatta gagactctgt tcaagctgga gcaacagatt 420
gtagcttctg gtggaatgat cccggccgtc gaactcggat tttga 465

```

<210> 22

<211> 817

<212> DNA

<213> Arabidopsis thaliana

<400> 22

```

gaaagacaaa acacaagttt cttcttcttg ggattggcta tttccagaaa tccaagtcaa 60
taatcaaagt ccaaacaaaa aaatcctctc ccaatctccg cttactctt ctcatggacg 120
ctctcattgc tcagcttcag agacaatttc gtgattacac catttctctc taccaacagg 180
ggtttttggg tgatcaattt actgagttga aaaagctaca agatgatgga agtcctgatt 240
ttgtgtctga agtgctttca ctttcttttg aagatttgtg gaagcttatc agtaacatgg 300
ctgaagcttt ggacacgaca ggaactgtag attttagtca ggtaggtgct agtgtgcatc 360
aattgaaggg tagtagctca agtgttggtg ccaagagggg caaaactttg tgtgttagct 420
tcaaggaatg ttgtgaagct aagaactacg aaggggtgtg gagatgtttg cagcaagtgg 480
atattgagta caaggcgta aagacaaagc tcaagatat gttcaatctt gagaaacaga 540
tcattcaagc tgggtggtata gttcctcaag tggatattaa ctaaagagac tagtccataa 600
gaagaaaaaa gatgatgact ttcttctttt agtttctctt cttaaattatt ttggatttgg 660
tgtttgcctc aaaactcaat aaaatatgtg caaaaagaaa caaaaacaag tgatggttgt 720
ttataaatca gtagtatgta ttgtttgatc tcatccgaga aaattgaaac cattggacta 780
atgaatgtga tgataatata tattggtttg cttctcc 817

```

<210> 23

<211> 510

<212> DNA

<213> Arabidopsis thaliana

<400> 23

```

aacttttagct atgaacacca tcgtcgttgc tcagttgcag agacaatttc aagactacat 60
cgtttctctt tatcaacagg gatttctgga taatcagttc tcagagttga gaaagttgca 120
agatgaagga acccctgatt ttgtagctga agttgtctct ctattcttcg acgactgttc 180
caagcttatt aataccatgt ctatatccct ggagcggcca gataatgtgg atttcaaaca 240
ggtggattca ggtgttcac aactcaaggg tagtagctcc agtgcgggtg caaggagggg 300
gaaaaatgtg tgcatactt tcaaggaatg ttgcgatgtt cagaaccgtg aaggggtgtc 360
aaggtgttta cagcaggtgg attatgaata taagatgtta aagactaaac ttcaggatct 420
ctttaattta gagaaacaga tcctccaagc tggaggtaca attcctcaag tggatataaa 480
ttagaccgat gcgtttctcg attatgcaaa 510

```

<210> 24

<211> 465

<212> DNA

<213> Artificial Sequence

<220>

<223> derived from Arabidopsis thaliana

<400> 24

```

atggatttgg ttcagaagca gaagagtttg caagattaca ccaaactact cttcttagaa 60
gggatttggg acagccagtt cttgcagctg caacaactac aagatgaaag caatccagat 120
tttgtttcac aagttgtcac actcttcttc caagactctg ataggattct caatgatctc 180

```


Asp Gly Tyr Val Ser Lys Pro Phe Glu Glu Glu Val Leu Tyr Thr Ala
 1155 1160 1165
 Val Ala Arg Phe Phe Glu Pro Cys
 1170 1175

<210> 27
 <211> 1036
 <212> PRT
 <213> Arabidopsis thaliana

<400> 27
 Met Ser Leu Phe His Val Leu Gly Phe Gly Val Lys Ile Gly His Leu
 1 5 10 15
 Phe Trp Met Leu Cys Cys Trp Phe Val Ser Trp Phe Val Asp Asn Gly
 20 25 30
 Ile Glu Asp Lys Ser Gly Leu Leu Val Gly Ser Val Gly Asp Leu Glu
 35 40 45
 Lys Thr Lys Met Thr Thr Leu Lys Lys Lys Asn Lys Met Trp Phe Trp
 50 55 60
 Asn Lys Ile Ser Ser Ser Gly Leu Lys Ile Pro Ser Phe Ser Tyr Gln
 65 70 75 80
 Phe Leu Gly Ser Val Lys Phe Asn Lys Ala Trp Trp Arg Lys Leu Val
 85 90 95
 Val Val Trp Val Val Phe Trp Val Leu Val Ser Ile Trp Thr Phe Trp
 100 105 110
 Tyr Phe Ser Ser Gln Ala Met Glu Lys Arg Lys Glu Thr Leu Ala Ser
 115 120 125
 Met Cys Asp Glu Arg Ala Arg Met Leu Gln Asp Gln Phe Asn Val Ser
 130 135 140
 Met Asn His Val Gln Ala Met Ser Ile Leu Ile Ser Thr Phe His His
 145 150 155 160
 Gly Lys Ile Pro Ser Ala Ile Asp Gln Arg Thr Phe Ser Glu Tyr Thr
 165 170 175
 Asp Arg Thr Ser Phe Glu Arg Pro Leu Thr Ser Gly Val Ala Tyr Ala
 180 185 190
 Met Arg Val Leu His Ser Glu Arg Glu Glu Phe Glu Arg Gln Gln Gly
 195 200 205
 Trp Thr Ile Arg Lys Met Tyr Ser Leu Glu Gln Asn Pro Val His Lys
 210 215 220
 Asp Asp Tyr Asp Leu Glu Ala Leu Glu Pro Ser Pro Val Gln Glu Glu
 225 230 235 240
 Tyr Ala Pro Val Ile Phe Ala Gln Asp Thr Val Ser His Val Val Ser
 245 250 255
 Leu Asp Met Leu Ser Gly Lys Glu Asp Arg Glu Asn Val Leu Arg Ala
 260 265 270
 Arg Ser Ser Gly Lys Gly Val Leu Thr Ala Pro Phe Pro Leu Ile Lys
 275 280 285
 Thr Asn Arg Leu Gly Val Ile Leu Thr Phe Ala Val Tyr Lys Arg Asp
 290 295 300
 Leu Pro Ser Asn Ala Thr Pro Lys Glu Arg Ile Glu Ala Thr Asn Gly
 305 310 315 320
 Tyr Leu Gly Gly Val Phe Asp Ile Glu Ser Leu Val Glu Asn Leu Leu
 325 330 335
 Gln Gln Leu Ala Ser Lys Gln Thr Ile Leu Val Asn Val Tyr Asp Ile
 340 345 350
 Thr Asn His Ser Gln Pro Ile Ser Met Tyr Gly Thr Asn Val Ser Ala
 355 360 365

Ser Ile Ser Gln Ala Leu Cys Thr Gly Ile Asp Pro Pro Ile Val Ile
 835 840 845
 Val Lys Pro Leu Arg Ala Ser Met Leu Ala Thr Leu Gln Arg Gly
 850 855 860
 Leu Gly Ile Gly Ile Arg Gln Pro Gln His Lys Gly Pro Pro Ala
 865 870 875
 Leu Ile Leu Arg Asn Leu Leu Leu Gly Arg Lys Ile Leu Ile Val Asp
 880 885 890 895
 Asp Asn Asn Val Asn Leu Arg Val Ala Gly Ala Leu Lys Lys Tyr
 900 905 910
 Gly Ala Asp Val Val Cys Ala Gln Ser Gly Ile Lys Ala Ile Ser Leu
 915 920 925
 Leu Lys Pro Pro His Gln Phe Asp Ala Cys Phe Met Asp Ile Gln Met
 930 935 940
 Pro Gln Met Asp Gly Phe Gln Ala Thr Arg Arg Ile Arg Asp Met Gln
 945 950 955
 Gln Gln Met Asn Lys Arg Ile Lys Asn Gly Gln Ala Leu Ile Val Gln
 960 965 970 975
 Asn Gly Asn Lys Thr Ser Trp His Leu Pro Val Leu Ala Met Thr Ala
 980 985 990 995
 Asp Val Ile Gln Ala Thr His Gln Cys Leu Lys Cys Gly Met Asp
 1000 1005 1010
 Gly Tyr Val Ser Lys Pro Phe Gln Ala Gln Gln Leu Tyr Arg Gln Val
 1010 1015 1020
 Ser Arg Phe Phe Asn Ser Pro Ser Asp Thr Gln Ser
 1025 1030 1035

<210> 28
 <211> 1057
 <212> PRT
 <213> Arabidopsis thaliana
 <400> 28

Met Asn Trp Ala Leu Asn Asn His Gln Gln Gln Gln Gln Gln Pro Arg
 1 5 10 15
 Arg Ile Gln Ile Ser Asp Ser Gln Ser Leu Gln Asn Leu Lys Ser Ser
 20 25 30
 Asp Phe Tyr Gln Leu Gly Gly Gly Ala Leu Asn Ser Ser Gln Lys
 35 40 45
 Pro Arg Lys Ile Asp Phe Trp Arg Ser Gly Leu Met Gly Phe Ala Lys
 50 55 60
 Met Gln Gln Gln Gln Gln Gln His Ser Val Ala Val Lys Met Asn
 65 70 75 80
 Asn Asn Asn Asn Asn Asp Leu Met Gly Asn Lys Lys Gly Ser Thr Phe
 85 90 95
 Ile Gln Gln His Arg Ala Leu Leu Pro Lys Ala Leu Ile Leu Trp Ile
 100 105 110
 Ile Ile Val Gly Phe Ile Ser Ser Gly Ile Tyr Gln Trp Met Asp Asp
 115 120 125
 Ala Asn Lys Ile Arg Arg Gln Gln Val Leu Val Ser Met Cys Asp Gln
 130 135 140
 Arg Ala Arg Met Leu Gln Asp Gln Phe Ser Val Ser Val Asn His Val
 145 150 155 160
 His Ala Leu Ala Ile Leu Val Ser Thr Phe His Tyr His Lys Asn Pro
 165 170 175 180
 Ser Ala Ile Asp Gln Gln Thr Phe Ala Gln Tyr Thr Ala Arg Thr Ala
 185 190


```

Ser Ser Asn Val Arg Leu Met Val Ser Ile Glu Asp Thr Gly Ile Gly
      660      665      670
Ile Pro Leu Val Ala Gln Gly Arg Val Phe Met Pro Phe Met Gln Ala
      675      680      685
Asp Ser Ser Thr Ser Arg Asn Tyr Gly Gly Thr Gly Ile Gly Leu Ser
      690      695      700
Ile Ser Lys Cys Leu Val Glu Leu Met Arg Gly Gln Ile Asn Phe Ile
      705      710      715      720
Ser Arg Pro His Ile Gly Ser Thr Phe Trp Phe Thr Ala Val Leu Glu
      725      730      735
Lys Cys Asp Lys Cys Ser Ala Ile Asn His Met Lys Lys Pro Asn Val
      740      745      750
Glu His Leu Pro Ser Thr Phe Lys Gly Met Lys Ala Ile Val Val Asp
      755      760      765
Ala Lys Pro Val Arg Ala Ala Val Thr Arg Tyr His Met Lys Arg Leu
      770      775      780
Gly Ile Asn Val Asp Val Val Thr Ser Leu Lys Thr Ala Val Val Ala
      785      790      795      800
Ala Ala Ala Phe Glu Arg Asn Gly Ser Pro Leu Pro Thr Lys Pro Gln
      805      810      815
Leu Asp Met Ile Leu Val Glu Lys Asp Ser Trp Ile Ser Thr Glu Asp
      820      825      830
Asn Asp Ser Glu Ile Arg Leu Leu Asn Ser Arg Thr Asn Gly Asn Val
      835      840      845
His His Lys Ser Pro Lys Leu Ala Leu Phe Ala Thr Asn Ile Thr Asn
      850      855      860
Ser Glu Phe Asp Arg Ala Lys Ser Ala Gly Phe Ala Asp Thr Val Ile
      865      870      875      880
Met Lys Pro Leu Arg Ala Ser Met Ile Gly Ala Cys Leu Gln Gln Val
      885      890      895
Leu Glu Leu Arg Lys Thr Arg Gln Gln His Pro Glu Gly Ser Ser Pro
      900      905      910
Ala Thr Leu Lys Ser Leu Leu Thr Gly Lys Lys Ile Leu Val Val Asp
      915      920      925
Asp Asn Ile Val Asn Arg Arg Val Ala Ala Gly Ala Leu Lys Lys Phe
      930      935      940
Gly Ala Glu Val Val Cys Ala Glu Ser Gly Gln Val Ala Leu Gly Leu
      945      950      955      960
Leu Gln Ile Pro His Thr Phe Asp Ala Cys Phe Met Asp Ile Gln Met
      965      970      975
Pro Gln Met Asp Gly Phe Glu Ala Thr Arg Gln Ile Arg Met Met Glu
      980      985      990
Lys Glu Thr Lys Glu Lys Thr Asn Leu Glu Trp His Leu Pro Ile Leu
      995      1000      1005
Ala Met Thr Ala Asp Val Ile His Ala Thr Tyr Glu Glu Cys Leu Lys
      1010      1015      1020
Ser Gly Met Asp Gly Tyr Val Ser Lys Pro Phe Glu Glu Glu Asn Leu
      1025      1030      1035      1040
Tyr Lys Ser Val Ala Lys Ser Phe Lys Pro Asn Pro Ile Ser Pro Ser
      1045      1050      1055
Ser

```

<210> 29

<211> 1057

<212> PRT

<213> Artificial Sequence

195	200	205
Asn Phe Glu Arg Glu Met	Phe Glu Arg Gln His	Asn Trp Val Ile Lys
210	215	220
Thr Met Asp Arg Gly Glu	Pro Ser Pro Val Arg	Asp Glu Tyr Ala Pro
225	230	235
Val Ile Phe Ser Gln Asp	Ser Val Ser Tyr Leu	Glu Ser Leu Asp Met
245	250	255
Met Ser Gly Glu Glu Asp	Arg Glu Asn Ile Leu	Arg Ala Arg Glu Thr
260	265	270
Gly Lys Ala Val Leu Thr	Ser Pro Phe Arg Leu	Leu Glu Thr His His
275	280	285
Leu Gly Val Val Leu Thr	Phe Pro Val Tyr Lys	Ser Ser Leu Pro Glu
290	295	300
Asn Pro Thr Val Glu Glu	Arg Ile Ala Ala Thr	Ala Gly Tyr Leu Gly
305	310	315
Gly Ala Phe Asp Val Glu	Ser Leu Val Glu Asn	Leu Leu Gly Gln Leu
325	330	335
Ala Gly Asn Gln Ala Ile	Val Val His Val Tyr	Asp Ile Thr Asn Ala
340	345	350
Ser Asp Pro Leu Val Met	Tyr Gly Asn Gln Asp	Glu Glu Ala Asp Arg
355	360	365
Ser Leu Ser His Glu Ser	Lys Leu Asp Phe Gly	Asp Pro Phe Arg Lys
370	375	380
His Lys Met Ile Cys Arg	Tyr His Gln Lys Ala	Pro Ile Pro Leu Asn
385	390	395
Val Leu Thr Thr Val Pro	Leu Phe Phe Ala Ile	Gly Phe Leu Val Gly
405	410	415
Tyr Ile Leu Tyr Gly Ala	Ala Met His Ile Val	Lys Val Glu Asp Asp
420	425	430
Phe His Glu Met Gln Glu	Leu Lys Val Arg Ala	Glu Ala Ala Asp Val
435	440	445
Ala Lys Ser Gln Phe Leu	Ala Thr Val Ser His	Glu Ile Arg Thr Pro
450	455	460
Met Asn Gly Ile Leu Gly	Met Leu Ala Met Leu	Leu Asp Thr Glu Leu
465	470	475
Ser Ser Thr Gln Arg Asp	Tyr Ala Gln Thr Ala	Gln Val Cys Gly Lys
485	490	495
Ala Leu Ile Ala Leu Ile	Asn Glu Val Leu Asp	Arg Ala Lys Ile Glu
500	505	510
Ala Gly Lys Leu Glu Leu	Glu Ser Val Pro Phe	Asp Ile Arg Ser Ile
515	520	525
Leu Asp Asp Val Leu Ser	Leu Phe Ser Glu Glu	Ser Arg Asn Lys Ser
530	535	540
Ile Glu Leu Ala Val Phe	Val Ser Asp Lys Val	Pro Glu Ile Val Lys
545	550	555
Gly Asp Ser Gly Arg Phe	Arg Gln Ile Ile Ile	Asn Leu Val Gly Asn
565	570	575
Ser Val Lys Phe Thr Glu	Lys Gly His Ile Phe	Val Lys Val His Leu
580	585	590
Ala Glu Gln Ser Lys Asp	Glu Ser Glu Pro Lys	Asn Ala Leu Asn Gly
595	600	605
Gly Val Ser Glu Glu Met	Ile Val Val Ser Lys	Gln Ser Ser Tyr Asn
610	615	620
Thr Leu Ser Gly Tyr Glu	Ala Ala Asp Gly Arg	Asn Ser Trp Asp Ser
625	630	635
Phe Lys His Leu Val Ser	Glu Glu Gln Ser Leu	Ser Glu Phe Asp Ile
645	650	655
Ser Ser Asn Val Arg Leu	Met Val Ser Ile Glu	Asp Thr Gly Ile Gly

